



Sensitivity analysis of InfoCrop model for Indian-mustard varieties in western Haryana

Yogesh Kumar*, Raj Singh, Anil Kumar and Sagar Kumar

Department of Agricultural Meteorology, CCS Haryana Agricultural University,
Hisar-125004, Haryana, India

* Corresponding author: yogeshgujjar62@gmail.com

(Received: 11 Jan 2017; Revised: 18 February 2017; Accepted: 13 May 2017)

Abstract

An InfoCrop model was validated with data sets generated on three varieties of *Brassica juncea* viz. Kranti, RH 406 and RH 0749 through field experiment laid in split plot design with three sowing dates of Rabi 2014-15 viz. 25th October, 5th November and 15th November, respectively. In sensitivity analysis, different combinations of maximum and minimum temperature, seasonal rainfall and elevated CO₂ concentration were simulated which reflected a substantial change in seed yield. A change of $\pm 1^{\circ}\text{C}$ in daily maximum and minimum temperatures led to increase in seed yield over base yield. The simulation results also supported the increase in seed yield in all three varieties due increase in the seasonal rainfall by 10 to 20%. Further, elevation in CO₂ concentration up to 490 ppm at constant temperature resulted in 13–32% also showed increment in seed yield of mustard; however, further increase in CO₂ concentration beyond 490 ppm adversely affected the seed yield all three varieties.

Key words: *Brassica juncea*, climate change, InfoCrop model, simulation

Introduction

Indian mustard [*Brassica juncea* (L.) Czern & Coss.] is much sensitive to climatic conditions and climate change could have substantial effects on its production. Global production of the annual crops is expected to reduce significantly due to climate change by the end of the 21st century. IPCC global studies indicate considerable probability of loss in crop production in India with increases in temperature (IPCC, 2014). According to fourth IPCC assessment report (AR₄), global average temperature has increased by 0.74 °C over the last hundred years and projected temperature increase is about 1.8 to 4 °C by 2100 A.D. Global losses may account from 1 to 5 % of GDP but, developing countries with tropical and sub-tropical climate are likely to suffer more (Parry *et al.*, 2007).

Studies on impact of climate change on crops need simulation models as they provide to quantify the effects of climate, soil and management on crop growth, productivity and sustainability of agricultural

production. These tools may reduce the expensive and time consuming field experimentations as they can be used to extrapolate the results of research conducted in one season or location to other season, location, or management (Boomiraj *et al.*, 2007). InfoCrop growth simulation model, developed under Indian condition, is one of the user friendly dynamic crop growth models. This model has the capability to estimate the potential yield and yield gaps, and also to assess the impacts of climate variability and climate change. This model simulates the crop growth processes viz. phenology, interaction among genotypes, environment, management and pests, yield forecast, yield loss assessment due to pests and greenhouse gas emissions (Aggarwal *et al.*, 2006). InfoCrop model can successfully simulate growth and yield of mustard crop across different locations in India. Simulated yield of mustard has been found to be sensitive to change in atmospheric CO₂ and temperature (Boomiraj *et al.*, 2010). An elevated level of CO₂ concentration and temperature shows negative impact on mustard yield (Singh *et al.*, 2013). Till now the scientific information on

simulation of growth and yield of mustard crop using modeling in Haryana state is very limited; therefore, keeping in view the importance of the role of crop simulation model in agriculture, the present investigation was carried out to validate and carry out sensitivity analysis of InfoCrop model for Indian-mustard crop under western Haryana conditions.

Materials and Methods

An experiment was designed to simulate the impact of change in maximum temperature (T_{max}) & minimum temperature (T_{min}), seasonal rainfall and

elevated CO_2 concentration within a range of $\pm 5^\circ C$, $\pm 10\%$ and 415 to 640 ppm, respectively, on the seed yield of three varieties of Indian mustard (*Brassica juncea*) viz. Kranti, RH 406, and RH 0749. An InfoCrop model (Aggarwal *et al.*, 2009) was used for the simulation study. The experiment was laid in split plot design in four replication at Research farms of Agril. Meteorology, CCSHAU, Hisar (Haryana) during *Rabi* 2014-15. The study material was sown at three different dates viz. 25th October, 5th November and 15th November, 2014. The meteorological data used in the present

Table 1: Simulated effect of change in daily (a) maximum and (b) minimum temperature on yield of mustard varieties

Change in	Variety						
	Kranti	RH 406	RH 0749	Kranti	RH 406	RH 0749	
maximum	Change in yield over base value (Kg ha ⁻¹)			% change			
temperature	5	1041.7	1213.0	1426.3	-21	-18	-16
	4	1120.8	1272.1	1460.2	-15	-14	-14
	3	1397.7	1612.3	1884.7	6	9	11
	2	1503.2	1686.3	1969.6	14	14	16
	1	1582.3	1878.6	2173.4	20	27	28
	-1	1661.4	1908.2	2190.3	26	29	29
	-2	1542.7	1715.9	2020.5	17	16	19
	-3	1199.9	1346.1	1562.1	-9	-9	-8
	-4	1041.8	1183.4	1409.3	-21	-20	-17
	-5	988.9	1153.8	1341.4	-25	-22	-21
Base yield (kg/ha)*1318.6		1479.2	1697.9	1318.6	1479.2	1697.9	
Change in	Variety						
	Kranti	RH 406	RH 0749	Kranti	RH 406	RH 0749	
minimum	Change in yield over base value (Kg ha ⁻¹)			% change			
temperature	5	1041.7	1213.0	1426.3	-21	-18	-16
	4	1120.8	1272.1	1460.2	-15	-14	-14
	3	1397.7	1612.3	1884.7	6	9	11
	2	1503.2	1686.3	1969.6	14	14	16
	1	1582.3	1878.6	2173.4	20	27	28
	-1	1595.5	1819.4	2088.5	21	23	23
	-2	1503.2	1701.1	1986.6	14	15	17
	-3	1226.3	1405.3	1646.9	-7	-5	-3
	-4	1147.2	1316.5	1545.1	-13	-11	-9
	-5	1081.2	1242.5	1443.2	-18	-16	-15
Base yield (kg/ha)*1318.6*		1479.2*	1697.9*	1318.6*	1479.2*	1697.9*	




Figure 1: Weekly weather parameter a) Temperature; b) Relative humidity; c) Rain, wind speed, SSH and PE during the crop season 2014-15 at Hisar

investigation was recorded at meteorological observatory of Department of Agricultural Meteorology during the standard weeks (SMW) from week 43rd to 15th and has presented in fig 1.

Results and Discussion Temperature

Maximum temperature: Simulated effect of varying levels of daily T_{max} on yield of mustard varieties has been detailed in table 1a and figure 2. The effect of both gradual increase and as well as decrease in the daily T_{max} (-5 to 5 °C) during the experimental period was studied in mustard. As a result, we recorded a change in mean yield, in a particular fashion, with respect to change in daily T_{max} (Figure 2). The yield was found to be increased when the temperature variation was kept between -2 to 3°C, however, the maximum yield was obtained at the T_{max} -1. Further, when the T_{max} was lowered and increase by -5 to -3 and 4 to 5, respectively, the yield loss was high. Substantial results were obtained with variation in between $\pm 1^\circ\text{C}$. The effect of percent change in yield was higher in RH 0749 followed by RH 406 and Kranti.




Figure 2: Depicting the InfoCrop simulation results of impact of change in (a) maximum temperature (T_{max}) (b) minimum temperature (T_{min}) (c) seasonal rainfall and (d) elevation in CO_2 concentration on the seed yield of all three varieties of mustard during the Rabi- 2014-2015

Table 2: Simulated effect of varying levels of rainfall on yield of mustard varieties

Per cent change in rainfall	Variety			Variety		
	Kranti	RH 406	RH 0749	Kranti	RH 406	RH 0749
	Change in yield over the base value (Kg ha ⁻¹)			% change		
50	1015.30	1668.58	1375.32	-23	-21	-19
40	1155.07	1298.75	1528.14	-12.4	-12.2	-10
30	1304.07	1421.52	1638.50	-1.1	-3.9	-3.5
20	1503.17	1698.13	2000.16	14	14.8	17.8
10	1402.96	1576.84	1850.74	6.4	6.6	9
-10	1354.17	1511.75	1755.66	2.7	2.2	3.4
-20	1305.38	1455.54	1657.18	-1	-1.6	-2.4
-30	1235.50	1377.15	1582.47	-6.3	-6.9	-6.8
-40	1124.74	1301.71	1511.16	-14.7	-12	-11
-50	962.56	1094.62.5	1205.53	-27	-26	-29
Base yield (kg/ha)*	1318.57*	1479.21*	1697.93*	1318.57*	1479.21*	1697.93*

Table 3: Simulated effect of varying levels of CO₂ on yield of mustard varieties

Change in CO ₂ concentration (ppm)	Variety			Variety		
	Kranti	RH 406	RH 0749	Kranti	RH 406	RH 0749
	Change in yield over base value (Kg ha ⁻¹)			% change		
415	1503.2	1671.5	1969.6	14	13	16
440	1582.3	1789.8	2071.5	20	21	22
465	1674.6	1849.0	2139.4	27	25	26
490	1740.5	1952.6	2224.3	32	32	31
515	1410.9	1568.0	1799.8	7	6	6
540	1213.1	1375.7	1579.1	-8	-7	-7
565	1094.4	1257.3	1477.2	-17	-15	-13
590	1015.3	1153.8	1324.4	-23	-22	-22
615	883.4	1020.7	1188.6	-33	-31	-30
640	777.9	887.5	1052.7	-41	-40	-38
Base yield (kg/ha)*	1318.6*	1479.2*	1697.9*	1318.6*	1479.2*	1697.9*

The simulated results indicates that decrease in T_{max} is more beneficial as compare to increase since elevated temperature lowers the days to flowering and days to maturity, which in turn lowers the total crop duration. An increase in daily T_{max} resulted in increased mustard yield which may be due to the pre proposed fact, that, in plants, warmer temperature accelerates growth and development leading to less time for carbon fixation and biomass accumulation before seed set resulting in poor yield (Rawson, 1992; Morison, 1996). Similar results were supported by

earlier studies of same aspect (Singh *et al.*, 2008; Kumar *et al.*, 2010; Aggarwal *et al.*, 2006).

Minimum temperature: The effect of change in daily T_{min} between -5 and 5°C was also studied for its impact on mustard yield (table 1b and figure 3). Interestingly, the impact on mustard yield, of change in T_{min} was quite similar to the impact of T_{max} . However, unlike T_{max} , here the highest benefits on yield was obtained on increasing the daily T_{min} by 1. The effect of percent change in the base yield was

increased by 1°C. Our results are in support with Singh *et al.* (2008) and Kumar *et al.* (2010).

Rainfall

In the present investigation substantial impact of change in seasonal rainfall in a range of $\pm 10\%$ on the seed yield of mustard was recorded. The details of simulated effect of varying levels of rainfall on seed yield of mustard are given in table 2 and figure 2(c). An acceptable degree of agreement was simulated by the InfoCrop simulation model under the elevated seasonal rainfall by 10 to 20%. Further, the pursuance of table 2 reveals that elevation in seasonal rainfall by 20% simulate the base yield at maximum level (RH 0749 > RH 406 > Kranti); However, under decreasing amount of seasonal rainfall varying from -10 to -50% model was under performed and gave the highest seed yield reduction in case of RH 0749 variety followed by Kranti and RH 406. Similar results were reported by earlier workers Pidgeon *et al.* (2001) and Singh *et al.* (2008).

CO₂ concentration

Simulated effect of varying levels of CO₂ on yield of mustard varieties are detailed in table 2 and figure 2(d). The InfoCrop also helped in simulating the effect of elevated CO₂ level (415 to 640 ppm) on the seed yield in the crop under study. The simulation showed positive impact on seed yield when CO₂ concentration was elevated up to 490 ppm. The further elevation in CO₂ concentration left negative impact on seed yield might be due to warming effect over the mean positive seed yield response for the mustard crop. Model also simulated the acceptable limit of elevated CO₂ which may be due to increased photosynthetic activity during the growth period and resulted in quit increasing trend with the specific plant geometric parameters such as leaf area, leaf weight, biomass production and numbers of grain. Our findings are supported by earlier workers Mishra *et al.* (1999), Uperty *et al.* (2003), Rotter and Van de Geijn (1999)

Conclusion

In the present investigation this model helped us in simulating the impact of varying levels of maximum and minimum temperature, seasonal rainfall and CO₂ on three varieties of mustard *viz.* Kranti, RH 406

and RH 0749. Change in T_{max} and T_{min} by -1 to 1°C, rainfall by 10 to 20% and CO₂ between 415 to 490 ppm showed beneficial impact on the crop and increased the seed yield considerably. The InfoCrop model was found useful tool in simulating the effect of various meteorological parameters on mustard and may help us in increasing the overall yield of mustard under continuously changing climatic conditions to fight for the issue of future food security at both national and international level.

References

- Aggarwal PK, Kalra N, Chander S and Pathak H. 2006. InfoCrop: A dynamic simulation model for the assessment of crop yields, losses due to pests, and environmental impact of agro-ecosystems in tropical environments. I. Performance of the model. *Agric Syst* **89**: 47-67.
- Aggarwal PK. 2009. Global Climate change and Indian agriculture; Case studies from ICAR.
- Boomiraj K, Chakrabarti B, Aggarwal PK, Choudhary R and Chander S. 2010. Assessing the vulnerability of Indian mustard to climate change. *Agric Eco Environ* **138**: 265-273.
- Boomiraj K, Chakrabarti B, Aggarwal PK, Choudhary R and Chander S. 2007. Impact of Climate Change on Indian mustard (*Brassica juncea*) in contrasting Agro-environments of the tropics. ISPRS Archives XXXVIII-8/W3 Workshop Proceedings: Impact of Climate Change on Agriculture.
- Parry ML, Canziani OF, Paultikof JP, van der Linden PJ and Hanon CE. 2007. Climate Change – Impacts, Adaptation and Vulnerability Technical Summary of Working Group II to Fourth Assessment Report of Intergovernmental Panel on Climate Change (eds), Cambridge University Press, UK, pp. 23–78.
- Pidgeon JD, Werker AR, Jaggard KW, Richter GM, Lister DH and Jones PD. 2001. Climatic impact on the productivity of sugar beet in Europe, 1961-1995. *Agril Forest Meteorol* **109**: 27-37.
- IPCC 2014. The Synthesis Report of the Intergovernmental Panel on Climate Change WG II: Impacts, vulnerability and adaptation.

- Kumar G, Chakravarty NVK, Kurothe RS, Sena DR, Tripathi KP, Adak T, Haldar D and Anuranjan. 2010. Effect of projected climate change on mustard (*Brassica juncea*). *J Agromet* **12**: 168-173.
- Mishra RS, Abdin MZ and Uprety DC. 1999. Interactive effects of elevated CO₂ and moisture stress on the photosynthesis, water relation and growth of *Brassica* species. *Crop Sci* **182**: 223-230.
- Morison JIL. 1996. Global environmental change impacts on crop growth and production in Europe. Implications of global environmental change for crops in Europe. *Asp Appl Biol* **45**: 62-74.
- Rawson HM. 1992. Plant responses to temperature under conditions of elevated CO₂. *Aust J Bot* **40**: 473-490.
- Rotter R and Van de Geijn SC. 1999. Climate change effects on plant growth, crop yield and livestock. *Climatic Change* **43**: 651-681.
- Singh M, Kalra N, Chakraborty D, Kamble K, Barman D, Saha S, Mittal RB and Pandey S. 2008. Biophysical and socioeconomic characterization of a water-stressed area and simulating Agri-production estimates and land use planning under normal and extreme climatic events: a case study. *Environ Monit Assess* **142**: 97-108.
- Singh SD, Chakrabarti B, Muralikrishna KS, Chaturvedi AK, Kumar V, Mishra S and Harit R. 2013. Yield response of important field crops to elevated air temperature and CO₂ level. *Indian J Agril Sci* **83**: 1009-1012.
- Uprety DC, Dwivedi N, Jain V, Mohan R, Saxena JM and Paswan G. 2003. Response of rice cultivars to the elevated CO₂. *Biol Plantarum* **46**: 35-39.