



Research Paper

Development of enzyme immunoassay for quantitation of mycelial biomass in *Brassica* genotypes against *Alternaria* blight

Antara Das, Mohd Tasleem, Manoj Singh, Dinesh Pandey, Anil Kumar and Gohar Taj*

Department of Molecular Biology and Genetic Engineering, G B Pant University of Agriculture and Technology, Pantnagar-263145, India

*Corresponding author: gohartajkhan@rediffmail.com

(Received: 12 Oct 2015; Revised: 5 Nov 2015; Accepted: 30 Dec 2015)

Abstract

Brassica crops faces a major infestation of *Alternaria brassicae*, a causal agent of *Alternaria* blight. Adopted chemical methods for controlling *Alternaria* infestation are ineffective and environmentally challenging. To bypass these challenges, strategies of enhancing innate immunity of *Brassica* plant resistance to pathogen, or suppression of virulence seems promising. For suppression of virulence early detection of pathogen, before symptom appearance is the main limitation. Efficient disease management appropriately buffered with resistance disease programme depends largely on the reliable identification and characterization of fungal isolate. Therefore, there is a need to develop a potential tool for immunological characterization of isolates of *A. brassicae*. Polyclonal antibody against total mycelia of *A. brassicae* was generated to develop ELISA based method for early detection and quantitation of mycelial biomass. Total protein was extracted from pathogen inoculated *Brassica* leaves (Varuna and PAB 9511) during different time interval of infection. The quantitative estimation of mycelial biomass of *A. brassicae* was estimated by indirect ELISA using polyclonal antiserum. When compared with healthy control, pathogen was not detected in the susceptible cultivar Varuna until after 6 h of post-inoculation. It was also observed that among the inoculated genotypes of *Brassica*, mycelium biomass was increased with disease progression in susceptible cultivar (Varuna), while the tolerant cultivar (PAB 9511) contained less. This demonstrated the efficacy of adopted immunological technique as a potent tool for detection, quantitation of mycelial biomass for characterization of differential tolerance against *Alternaria* blight in *Brassica* genotypes.

Key words: *Alternaria brassicae*, *Alternaria* blight, ELISA

Introduction

Rape seed mustard is the second most edible oil seed crop in India. Compare to other edible oils, the rape seed mustard oil has the lowest amount of harmful saturated fatty acids and it also contains adequate amount of the two essential fatty acids linoleic and linolenic acid. Productivity of rapeseed mustard in India varies significantly from state to state, depending on the climatic conditions and other environmental factors prevailing in the respective areas. Low productivity of rape seed is attributed to its susceptibility of various diseases and the major one is *Alternaria* Blight (Meena *et al.*, 2010). A special pattern of development of several *Alternaria* species

on leaf discs of host and non host is known. When pathogen attacks on leaf, firstly the germination of spore takes place followed by germ tube formation and formation of appressorium. It is found that in host and non host species pathogen penetration is not different. However, plant responses to attempt penetration, which includes the formation of callose-containing papillae, callose deposition in the walls of attacked cells and their neighbouring cells and cell necrosis, varies with specific pathogen–plant interactions (McRoberts *et al.*, 1996). Pathogen activities focus on colonization of the host and utilization of its resources, while plants are adapted to detect the presence of pathogens and to respond

with antimicrobial defenses and other stress responses (Jinrong *et al.*, 2002). Early detection and correct identification of fungal pathogens is of primary importance in determining the most effective course of treatment to prevent the spread of fungi causing plant disease and post harvest storage losses. Thus in many aspects of plant pathology, there is a need for assay systems such as immunoassays that are specific, sensitive and quick to confirm visual systems, to detect early and latent infections before symptoms are visible (Dewey *et al.*, 1989; Xia *et al.*, 1992; Brill *et al.*, 1994). Most of the antibodies in fungal antisera that are non specific are of the immunoglobulin class IgM, whereas many of the antibodies that are species specific are IgG antibodies (Cahil and Hardham 1994; Thronton *et al.*, 1993, 1994). Use of such highly specific antibodies can help in identification of pathogen races and thus can help in pathotyping (Mohan, 1988). ELISA has been proven to be useful for monitoring the antigenic properties in the process of population shift during the sexual development of fungus together with the determination of genetic diversity among the pathogens and subsequently for immunopathotyping. Using *Stachybotrys chartarum* as a model fungus Schmechel *et al.* (2006) demonstrated that monoclonal antibodies (mAbs) can provide species specific diagnostic reagents and also be used to investigate immunological cross-reactivity patterns among fungi. They demonstrated that mAbs and immunoassays have the potential to detect *S. chartarum* species-specifically. Kamraj *et al.* (2012) studied on *Aspergillus oryzae*, extracted protein from the fungi and used it as an antigen to developed probe. This probe was used to develop a sensitive, rapid, and inexpensive screening test to detect *Aspergillus* in agricultural commodities. Dewey *et al.* (1990) studied on *Penicillium islandicum*, a fungus commonly isolated from stored rice grains in South-East Asia. They raised monoclonal antibodies (mAbs) against *Penicillium islandicum* and hybridoma supernatants were screened by ELISA. Most of the mAbs raised cross-reacted with other storage fungi or uninfected rice grains but three were species-specific. One of these, PIO1, was used to develop ELISA and DIP-STICK assays for the detection of *P. islandicum* in individual grains. Fox

et al., (2004) also developed an indirect immunofluorescence spore assay (IFSA) to detect urediniospores of *Phakopsora pachyrhiz*. Shamala *et al.* (2006) studied that mAbs raised against *G. boninense* were not specific as the mAbs gave positive signals through the cross-reactivity test with all fungi tested in the cross-reactivity. Schmechel *et al.* (2008) studied on *A. alternata*. They developed pAbs (polyclonal antibodies) against *A. alternata* which showing cross-react broadly with related and nonrelated fungi.

In our study polyclonal anti-mycelial antibodies is developed against mycelia antigen(s) of *A. brassicae*. Concentration of mycelial antigen(s) and antimycelial antibodies were optimized for development of enzyme immunoassay. Cross reactivity studies was conducted with different isolates of *A. brassicae*.

Materials and Methods

Source of plant material

Seeds of two *B. juncea* cultivars one PAB 9511 (tolerant) and the other Varuna (susceptible) used in this study were obtained from CRC Pantnagar, India and grown in pots containing mixtures of soil: sand: vermicompost (2:1:1). Plants were kept in a glass house at 22°C for 16 h photoperiod.

Isolation of *Alternaria brassicae* isolates

Ten genotypes of *A. brassicae* were taken, viz Ashirwad (AS), Divya (DV), JD6 (JD), Kanti (KA), Kranti (KR), Local Mustard (LM), NDRE4 (ND), PRE2007 (P7), PRE2010 (P10), Varuna (VR) obtained from Crop Research Centre, Pantnagar, India. Isolates were grown on potato dextrose broth media and incubated at 22°C in the dark for the 15 days to allow growth of the pathogen.

Artificial inoculation of pathogen

At the four leaf stage, plants of PAB 9511 and Varuna cultivars were inoculated with spore suspension (10^5 per ml) of *A. brassicae*. Fungal Spores were taken from infected leaves of *B. juncea* cultivar Varuna. *A. brassicae* inoculated plants were placed in moist chamber in a greenhouse at 22°C and 100% of relative humidity (RH). Control treatment was treated with sterilized distilled water.

Serological studies

Preparation of antigen

The antigen was prepared from total protein extracted from mycelium of *A. brassicae* isolate NDRE4. The mycelial mat was filtered through muslin cloth, washed with phosphate buffer saline (pH 7.4) and lyophilized. Lyophilized mycelium (1 mg) was ground with liquid N₂. PBS was added to the ground mycelium (El-Nashaar *et al.*, 1986) and the suspension was centrifuged at 12000 rpm for 30 min at 4°C. The supernatant containing total protein was obtained and protein concentration was determined by Bradford estimation (Bradford, 1976). Total protein concentration was adjusted to 1 mg mL⁻¹ and stored at -20°C.

Preparation of Antimycelial antibody

Antiserum against *A. brassicae* was prepared using New Zealand white rabbit by four weekly subcutaneous injections with mycelial antigen of 1 mg mL⁻¹. First dose was given by antigen emulsified with an equal volume of Freund's complete adjuvant (Difco lab, USA) and for subsequent boosting doses were used by Freund's incomplete adjuvant. The doses were administered at one week interval. Two weeks after the last injection, rabbits were bled and the blood was collected from the marginal ear vein and left for two hours at room temperature for clot formation then kept in refrigerator overnight. The antiserum was clarified by centrifugation at 3000g for 30 min at 4°C and stored at -20°C until use (Srivastava and Arora, 1997).

ELISA for detection of changes in antigenic determinants

The ELISA was developed with minor modification as described initially by Engvall and Perlmann, (1971) and later by Clark and Adams (1977). Lyophilized mycelium was dissolved in PBS at the rate of 20 mg/ml. Polystyrene micro plates were coated with mycelia fragments and plant extract treated mycelium harvested at 14th day and allowed incubating overnight at 4°C. The plates were washed with PBS+Tween-20 (0.01%). The plates were filled with PBS containing 5% skimmed milk for 2hr. at RT. 100µl (1:250 dilution) of 1st antibody

(Raised against respective antigens) were added and incubated for 2 hrs at RT. Washing was done thrice with PBS + Tween-20 (0.01%) + 0.25% BSA. 100µl of 1:1000 diluted alkaline phosphate conjugated secondary antibody was incubated for 2hr at room temperature. plates washed with PBS containing 0.25% BSA + Tween-20 (0.01%) thrice. Alkaline phosphate activity was assayed with p-nitrophenyl phosphate sodium salt. The plates were incubated for 30 min in dark and the reaction was stopped with 100µl of 1.5 M NaOH solution. Reading was taken at 405nm in ELISA reader.

Results and Discussion

Immuno pathotyping

Polyclonal antibodies raised against whole lyophilized mycelia of virulent pathogen of *A. brassicae*. Antimycelial antibody was used for development of immunoassay systems, viz indirect Enzyme linked immunosorbent assay (ELISA) procedure for pathogen identification and characterization. The immunoassays were developed performing antigen concentration kinetics and antibody dilution curve analysis. These assays were employed for immunological based characterization of diversity among these isolates of pathogen by studying antibodies reactivity pattern of pathogen isolates.

Optimization of Antigen kinetics through ELISA

In order to determine the virulence level of *A. brassicae* isolate/strains, the causative agent of Alternaria blight disease of Brassica and differentiating it from similar, *A. brassicae* isolate, mycelial

Table S1: Tabular representation of antigen kinetics of fungal antigen

Mycelial antigen in ng	Absorbance OD at 405 nm
50	0.638
100	0.659
250	0.7545
500	0.608
1000	0.4515
2000	0.4595
4000	0.3835
8000	0.378




Figure 1: Antigen kinetics for determination of optimal mycelial antigen concentration of *A. brassicae* for development of immuno-assay using anti-mycelial antibodies

fragments antigen from all isolates were subjected to indirect ELISA, standardization of antigen concentration for mycelial antigen was performed on ELISA plates. According to the result obtained, It was concluded that 250 ng/well of antigen

concentration was appropriate for determining the virulence level of *A. brassicae* (fig. 1).

Determination of titre of Anti-mycelial antibody

To determine the optimal antibody dilution, the




Figure 2: Determination of titre of Anti-mycelial antibody using indirect ELISA

Table S2: Tabular representation of anti-mycelial antibodies dilution as indirect ELISA absorbance values (405 nm)

Anti-mycelial antibodies dilution	Absorbance OD at 405
1:100	0.946
1:250	0.873
1:500	0.842
1:1000	0.663
1:2500	0.480
1:5000	0.169
1:10000	0.146
1:20000	0.021

micro-titre plates were coated with optimal amount of mycelial antigen that falls within the detection range of indirect ELISA i.e. 250 ng per well. Antibody dilutions for anti-mycelial antibody raised was used ranging from 1:100 to 1:20,000. At dilution, 1:5,000, reactivity corresponding to O.D.⁴⁰⁵ was observed 0.169 for mycelial antigen of *A. brassicae*. It was observed that the straight line was obtained after the antibody dilutions of 1:5000 and almost

stagnant immuno reactivity in the antibody dilution range of 1:5000 to 1:20,000 indicating that dilutions higher than this will not improve the binding activity. Thus 1:5000 was determined as titre of anti-mycelial antibody raised against mycelial antigen. The titre of antibodies is defined as maximal dilution of antibodies at which reactivity was visible. It was concluded that 1:250 antibody dilution is appropriate for determine the virulence level of *A. brassicae* (fig 2).

Cross reactivity patterns of different isolates of *Alternaria brassicae* using indirect ELISA

In order to increase the sensitivity of the analysis, compares the ELISA reactivity of anti-mycelial antibody against 10 isolates of *A. brassicae* pathogen and *A. solani*. These intra-species and intra-generic pathogen were showed cross-reactivity with anti-mycelial antibody raised against whole mycelial antigen of *A. brassicae*. Antimycelial antibodies strongly crossed react with mycelium of *A. brassicae* isolate and *A. solani*. Results indicated that different fungi express different amount per unit biomass of the antigen-antibody




Figure 3: Cross reactivity studies between different strain of *Alternaria brassicae* by indirect ELISA assay using anti-mycelial antibody and Dilution of primary antibody is 1:250, Dilution of secondary antibody is 1:1000, Antigen concentration is 250ng in 100 µl.

Table S3: Tabular representation of cross reactivity of *A. brassicae* isolates.

<i>Alternaria brassicae</i> isolates	Average Absorbance OD at 405 nm
PRE-07	0.528
JD-6	0.443
KRANTI	0.476
LOCAL MUSTARD	0.443
MNRE-4	0.482
ASHIRWAD	0.441
DIVYA	0.462
PRE-10	0.489
VARUNA	0.571
<i>A. solani</i>	0.456

binding sites for anti-mycelial antibody (fig. 3).

Serological Detection of Alternaria blight during different time intervals/stages of infection of *Alternaria brassicae* in host species of *Brassica juncea*

Total protein was extracted from artificially inoculated *Brassica* leaves (varuna and PAB 9511) at 3 hpi, 1dpi, 2dpi and 3dpi). Total protein extracted from healthy *Brassica* leaves was used as control.

Table S4 : Serological test for *A. brassicae* antiserum as indirect ELISA absorbance values (405 nm) of total protein extracted from inoculated brassica leaves after different periods of inoculation with *A. brassicae*.

Brassica cultivar Time post-inoculation	Absorbance values (405 nm)
Varuna (Susceptible)	
CV	0.539333
6 h	0.646333
1 D	0.845333
2 D	1.07
3 D	0.889333
PAB 9511 (tolerance)	
6 h	0.676667
1 D	0.583333
2 D	0.501667
3 D	0.5

Indirect ELISA was used to detect *Alternaria* blight pathogen. *Alternaria brassicae* was used at a dilution of 1:250 and 1 µg protein per well in serological tests. The amount of mycelium of *A. brassicae* was estimated by indirect ELISA after different periods of inoculation with *A. brassicae* using polyclonal antiserum specific to *A. brassicae*, in leaves of tolerant (PAB9511) and




Figure 4: Serological Detection of *A. brassicae* in inoculated Brassica genotype of Varuna and PAB 9511 of different time interval

susceptible (varuna) *Brassica* cultivars, by comparing absorbance values obtained with those given by control samples for each isolate. Absorbance values of at least double of that of the healthy control were considered positive. Compared with healthy control, pathogen was not positively detected in the susceptible *Brassica* (varuna) until after 6 h post-inoculation in inoculated leaves (0.654). After that, a sharp increase in ELISA values was recorded with the time (fig-4). Low levels of ELISA value were obtained in case of tolerant *Brassica* cultivar (PAB 9511). The amount of antigen in leaves was sufficient to cause a positive ELISA reaction (0.853 to 1.46, respectively) only after 6 h post-inoculation. Results indicated that among the inoculated *Brassica* (6h to 3days post-inoculation), there was large amount of mycelium in leaves of the susceptible *Brassica* cultivar (Varuna), while the tolerant *Brassica* cultivar (PAB 9511) contained less. The experiment was repeated twice and ELISA absorbance values at 405 nm was average of two replicates each and 100 μ l from the diluted *A. brassicae* antiserum (1:250 diluted in PBS) were used for each ELISA reaction. Serological testing can be useful for early detection of the presence of *A. brassicae* before appearance of the symptoms. Keeping in view the present study was carried out to generate polyclonal antibody against total mycelia of *A. brassicae* to develop ELISA based method for early detection and quantitation of mycelia biomass. It could be possible that polyclonal antibodies raised against specific strain of *A. brassicae* were showing cross reactivity with other strain of *A. brassicae* and other species of *Alternaria*.

From the above study it was concluded that polyclonal antibodies against mycelial antigen of *A. brassicae* was showing the interaction with *A. brassicae*. Optimal concentration of mycelial antigen of *A. brassicae* was found to be 250 ng/well for development of indirect ELISA method for detection of *Alternaria* blight disease. It was concluded that 1:250 anti-mycelial antibody dilution was appropriate for anti-mycelial antibody to determine the virulence level of *A. brassicae*. The titer of anti-mycelial antibody generated against mycelial antigen was 1:5000. Cross reactivity studies was done by indirect ELISA. Due to polyclonal nature

of anti-mycelial antibody cross-reactivity pattern was observed among different isolates of mycelial antigen of *A. brassicae*. Serological detection of *A. brassicae* was done in inoculated *Brassica* genotype of varuna and PAB 9511 with different time interval. It was observed that among the inoculated *Brassica* (6h to 3days post-inoculation), the mycelial biomass was increased with disease progression. Gemmrich and Seidel (1996) found out the detection of *Oidium* mycelium by using polyclonal antiserum. In the present study, it was also observed that relatively more amount of mycelium biomass was observed in inoculated leaves of susceptible *Brassica* cultivar (varuna), as compared to tolerance *Brassica* cultivar (PAB 9511).

In this study it was observed that the polyclonal antibody was helpful to determine pathogen biomass in infected leaves. By using ELISA we can determine the level of infection. For future prospective it could be used for quantitation of mycelial biomass in plant patho-system. Kennedy *et al.* (1999) developed detection method of *Mycosphaerella brassicicola* in infected crop of vegetable brassicas. We need to work at the production of additional mAbs against the purified antigen of *A. brassicae*. This will allow the development of a mAb-based sandwich assay for environmental sample analysis. In conclusion, the widespread cross-reactivity among fungi and the observed biological variability are considered to be major challenges for the successful implementation of accurate and quantitative immunoassays.

Acknowledgements

Authors are grateful to the Department of Biotechnology (DBT), Government of India, New Delhi, India for providing financial assistance under HRD Program and project of Program Mode Support for Research and Development in Agricultural Biotechnology.

References

- Brill LM, McClary RD and Sinclair JB. 1994. Analysis of two ELISA formats and antigen preparations using polyclonal antibodies against *Phomopsis longicolla*. *Phytopathol* 84: 173-179.
- Bradford MM. 1976. A rapid and sensitive method for the quantitation of binding. *Analytical Biochem* 72: 248-254.
- Clark MF and Adams AN. 1977. Characteristics of the microplate methods of enzyme-linked

- immunosorbent assay for the detection of plant viruses. *J General Virol* **34**: 475-483.
- Cahill DM and Hardham AR. 1994. Exploitation of zoospore toxin in the development of novel clipstick immuneassay for the detection of *Phytophthora cinnamomolmi*. *Phytopathol* **84**: 193-200.
- Dewey FM, MacDonald MM and Phillips SI. 1989a. Monoclonal Antibodies to specific components on the dutch Elm pathogen *Ophiostoma ulmi*. *Plant Pathol* **38**: 9-20.
- Dewey FM, MacDonald MM and Phillips SI. 1990. Development of monoclonal antibody-ELISA, Dot-Blot and DIP-STICK immunoassay for *Penicillium islandicon* in rice grains. *J Gen Microbiol* **136**: 753-760.
- Dewey FM, Macdonald MM, Phillips SI and Priestley RA. 1990. Development of monoclonal-antibody-ELISA and -DIP-STICK. *J General Microbiol* **135**: 361-374.
- Engvall E and Perlmann P. 1971. Enzyme-linked immunosorbent assay (ELISA) quantitative assay of immunoglobulin G. *Immunochemistry* **8**: 871-874.
- Fox M, Gray G, Kavanagh K, Lewis C and Doyle S. 2004. Detection of *Aspergillus fumigatus* mycotoxins immunogen synthesis and immunoassay development, *J Microbiol Methods* **56**: 221- 230.
- Gemrich AR and Seidel M. 1996. Immunodetection of overwintering *Oidium* mycelium in bud scales of *Vitis vinifera*. *Vitis* **35**: 63-64
- Kamaraj M, Rajeshwari Sivaraj and Muthu Samygovarathan .2012. Raising of polyclonal antibody against the *Aspergillus oryzae* proteins in albino rat. *Inter J Pharma Bio Sci* **3**: 342-348.
- Kennedy R, Wakeham A J and Cullington J E. 1999. Production and immunodetection of ascospores of *Mycosphaerella brassicicola*: ringspot of vegetable crucifers, *Plant Pathology* **48**: 297-307.
- Jinrong Wan F, Andrew MD and Bent F. 2002. Probing plant-pathogen interactions and downstream defense signaling using DNA microarrays. *Funct Integr Genomics* **2**: 259-273.
- McRoberts N and Lennard JH. 1996. Pathogen behaviour and plant cell reactions in interactions between *Alternaria* species and leaves of host and nonhost plants. *Plant Pathol* **45**: 742-752.
- Meena PD, Awasthi RP, Chattopadhyay C, Kolte SJ and Kumar A. 2010. *Alternaria* blight: a chronic disease in rapeseed-mustard. *J Oilseed Brassica* **1**: 1-11
- Mohan SB. 1988. Evaluation of Antisera raised against *Phytophthora fragariae* for detection of the red core disease of strawberries by enzyme-linked immunosorbant assay (ELISA). *Plant Phytopathol* **37**: 206-216.
- Shamala S, Chris D, Sioban O and Idris A. 2006. Preliminary studies on the development of monoclonal antibodies against mycelia of *Ganoderma boninense*, the causal pathogen of basal stem rot of oil palm. *Mal J Microbiol* **2**: 30-34.
- Srivastava AK and Arora DK. 1997. Evaluation of a polyclonal antibody immunoassay for detection and quantification of *Macrophomina phaseolina*. *Plant Pathol* **46**: 785-794.
- Schmechel D, Simpson JP, Beezhold D and Lewis DM. 2006. The development of species-specific immunodiagnosics for *Stachybotrys chartarum*: The role of cross-reactivity. *J Immunol Methods* **309**: 150 – 159.
- Schmechel D, Gree BJ, Blachese FM, Janotka E and Beezhold DH. 2008. Analytical bias of croos-reactive polyclonal antibodies for environmental immunoassays for *Alternaria alternata*. *J Allergy Clin Immunol* **121**: 63-768.
- Thronton CR, Dewey FM and Gilligan CA. 1993. Development of monoclonal antibody based immunological assays for the detection of live propagules of *Rhizoctonia solani* in soil. *Plant Pathol* **42**: 763-773.
- Thronton CR, Dewey FM and Gilligan CA. 1994. Development of monoclonal antibody based enzyme-linked immunosorbant assay (ELISA) for the detection of live propagules of *Trichoderma harzianum* in a peat /bran medium. *Soil Biol Biochem* **26**: 909-920.
- Xia JQ, Lee FN, Scott HA and Raymond LR. 1992. Development of monoclonal antibodies specific for *Pycularia grisea*, the rice blast pathogen. *Mycological Res* **96**: 867-873.