



Pathogenicity test of *Alternaria brassicae* (Berk.) Sacc. using artificial inoculation methods on common varieties of rapeseed-mustard in Punjab region

Ram Singh Dhaliwal and Bahaderjeet Singh*

University college of Agriculture, Guru Kashi University, Talwandi Sabo, 151302

*Corresponding author: sidhubahaderjit@yahoo.in

(Received: 5 Dec 2018; Revised: 10 Dec 2018; Accepted: 20 Dec 2018)

Abstract

To develop the precision, sensitivity for pathogenicity testing, a comparison was carried out among three different artificial inoculation methods namely foliar spraying method, drop method and biocontrol+foliar spray method for pathogenesis studies. The pathogen *A. brassicae* was isolated and from infected mustard plants. The drop inoculation method was most ideal as this fixed the inoculum on the target site. Drop method produces significantly higher number of disease lesions than the other inoculation methods and has the advantage of being accurate and precise. The number of initial disease lesions in drop method was highest in all of the time intervals of observation namely 71 in *B. juncea* (Giriraj), 59 in *B. juncea* (RLC-3), 53 in *B. napus* (GSC-7), 73 in *B. nigra* (PC-6) at 120h after pathogen inoculation compared to other methods respectively. The results in this particular study indicates that two varieties *B. juncea* (RLC-3) and *B. napus* (GSC-7) showed some tolerance as compared to other two varieties *B. juncea* (Giriraj) and *B. nigra* (PC-6). The biocontrol agent used in particular study showed that the tolerance can be induced in the plants using *Trichoderma viride*. As the number of lesions and pathogenicity caused by *A. brassicae* was reduced considerably when the culture of *T. viride* was used.

Key words: *Alternaria brassicae*, artificial inoculation, *Brassica* spp., slide culture technique, *Trichoderma* spp.

Introduction

Oilseed brassicas, also known as rapeseed-mustard, constitute the second largest group of Oilseed crops that contributes 32 per cent of the total Oilseed production in India. In India rapeseed mustard occupy an area of 57.62 lac ha with a production of 68.22 lac tons and productivity of 1184 kg / ha (Anon, 2016). Brassica oilseed crops play an important role in the diversification in cropping system and also in providing the quality food by meeting the fat requirement to same extent. However, the productivity of these crops is relatively low, thus there is a significant shortfall in the supply of required quantity of fat per capita (presently available of required per capita), no doubt there are several other crops like groundnut, sunflower etc. therefore huge amount of foreign exchange is drained for imported the huge quantity of oilseeds to meet the national requirement.

In Punjab, the area under brassica oilseed is decreasing under rapeseed mustard for the last 10years. This is due to the fact that the alternate Rabi crop of wheat is more profitable as the brassica crops are more vulnerable to several biotic and abiotic stresses. Of these the attack of diseases like *Alternaria* blight is more concerning point it is most destructive and wide spread affecting all the cultivated species. Highest yield obtained from a

particular crop and averages realized at the state and national levels. A major contributory factor to this gap is its unchallenged exposure to a number of biotic, mesobiotic and abiotic stresses. This disease is also called the black spot (Louvét, 1958) or grey spot (McDonald, 1959) based on the symptoms produced on the host. *Alternaria* blight causing pathogens can attack all the aerial parts of plant and can cause huge losses in yield. *Alternaria* blight caused by most *Alternaria brassicae* (Berk.) Sacc. has been reported from all the continents of the world affecting cruciferous crops. The disease appears annually during the cropping season (from October to February) in different parts of India and causes enormous loss to growers (Prasad *et al.*, 2006). Environmental factors play an important role in the development of the disease. The environmental variables viz., temperature, humidity, rainfall and sunshine are the most critical factors. They affect the pathogen and host or host-pathogen interaction during pathogenesis.

The ideal and most economical mean of managing the *Alternaria* blight disease of rapeseed-mustard would be the use of resistant varieties. Most of the cost effective and eco-friendly management strategies for evolving selecting the genotypes possessing the resistant/tolerance reaction against the diseases. Information on

resistance source is not available, although some sort of tolerance may be available (Shah *et al.*, 2005). Biocontrol agents can be effective to induce tolerance against *A. brassicae* and *A. brassicola*. Reshu and Khan (2012) reported *Trichoderma viride* isolated from mustard leaf showed fungicidal activity against *A. brassicae*. Pathogenicity testing using artificial inoculation is easy and less time consuming technique to study the various aspects of plant pathology, including epidemiology, etiology, disease resistance, host-parasite interaction and disease control (Xu and Ko, 1998).

Detached leaf inoculation method has been found to be the elementary and swift method. The conventional method like spraying has the disadvantage of causing considerable variation in spore distribution (Tuite, 1969). Although the accuracy and precision were improved by applying a drop of inoculum with a modified hypodermic needle (Lapwood and Mckee, 1966) capillary pipette (Toussoun *et al.*, 1960). In this report, to develop precision, sensitivity and for testing a comparison was carried out among three different artificial inoculation methods namely foliar spray method, drop method and biocontrol+ spray method for pathogenesis studies of *A. brassicae* on *B. juncea* (Indian mustard), *B. napus* (Gobhi Sarson) and *B. nigra* (Banarasi Rai).

Materials and Methods

Isolation: *A. brassicae* (Berk.) Sacc. was isolated from a diseased leaf of *B. juncea*, *B. napus* and *B. nigra* from Guru Kashi University Research Farm Talwandi Sabo, Bathinda (Punjab), India. The pathogen was isolated, purified by single spore technique (Toussoun and Nelson, 1976), and maintained on potato dextrose agar (PDA) media at $25^{\circ}\pm 1^{\circ}\text{C}$.

Identification using slide culture technique: The culture exhibiting maximum production was identified by both conventional methods (morphological and microscopic methods). Slide culturing was done to determine the microscopic features for morphological characterization according to standard taxonomic key (Singh *et al.* 2012; Singh *et al.* 2013.)

Inoculum preparation: Seven-day old *Alternaria* cultures grown on PDA plates were flooded with distilled water and spores were released by agitation with a sterile brush. The spores in this suspension were counted using a haemocytometer and the concentration was adjusted to 10^4 spores/ml.

Detached leaf inoculation: Leaves were detached from

the *B. juncea* (2 varieties- 'Giriraj' and 'RLC-3'), *B. napus* (1 variety- 'GSC-7') and *B. nigra* (1 variety- 'PC-6') washed under running tap water followed by a wash with autoclaved water and then surface was wiped off with 70% ethanol.

Pathogenicity test were carried out with foliar spray, drop method and biocontrol (*Trichoderma viride*) + spray method on detached leaves of *B. juncea*, *B. napus* and *B. nigra*. In the spray method, inoculum was sprayed on detached leaves with the help of atomizer. In drop method; inoculum was placed on each detached leaf in the form of drop. In Biocontrol+spray method, inoculum was sprayed on leaf followed by (*Trichoderma viride* 1.5% W.P) 1mg/ml suspension and go on with 10^4 spores/ml. Detached leaves were kept in sealed plastic bags with moistened filter paper and placed in BOD incubator at $25^{\circ}\pm 1^{\circ}\text{C}$ and 70% relative humidity. Wetness of filter paper was maintained by spraying autoclaved distilled water. The plastic bags were observed for *A. brassicae* initial symptoms at intervals of 24 h after pathogen inoculation and the infected leaves were examined up to 120 h after pathogen inoculation. The numbers of disease lesions were counted on the detached leaves in all the inoculation methods at 24, 48, 72, 96 and 120 h after inoculation of pathogen while the control remains free from severe symptoms.

Results and Discussion

The fungus was identified according to standard taxonomic key including colony colour, and the morphology of hyphae and conidia. Colonies were fast growing, black to olivaceous-black or greyish colour when incubated on PDA plates at 30°C for 7 days (Fig. 1a). Septate hyphae, conidiophore structure and development of conidia was seen under the microscope (Fig. 1b,c). Conidia were found in chains with a conspicuous beak, smooth, greyish-olive, with 7-8 cross septa and longitudinal ones, slightly constricted at the septa (Fig. 1d).

In this scrutiny, four varieties of *Brassica* spp. were taken as the criteria to find the numbers of lesions on the detached leaves in an incubation hours of 24, 48, 72, 96, 120 with different types of artificial inoculation methods namely spray, drop and biocontrol+spray methods. The best artificial inoculation method was analysed for the pathogenesis studies.

The *B. juncea* (1st variety- Giriraj) at 48 h after inoculation of pathogen, drop method showed highest number of initial lesions that is (39) lesions, followed by spray (15) lesions and biocontrol+spray (1). Maximum number of lesions i.e. 71, 37 and 3 were appear on leaves of *B. juncea* using drop, spray and biocontrol+spray method,




Fig.1. a) Colony morphology. b) Structure of conidiophore and development of conidia. c) Septate Hyphae. d) Chain of conidia.




Fig. 2. Number of lesions on the detached leaf of *B. juncea* (variety-Giriraj) using different pathogen inoculation methods (drop, spray and biocontrol+ spray) at different time intervals (24, 48, 72, 96 and 120 h) after pathogen inoculation.




Fig. 4. Number of lesions on the detached leaves of *B. juncea* (variety- RLC-3) using different pathogen inoculation methods (drop, spray and biocontrol+ spray) at different time intervals (24, 48, 72, 96 and 120 h) after pathogen inoculation.




Fig 3. Development of initial black/brown disease lesions on the detached leaves of *B. juncea* variety Giriraj using different pathogen inoculation methods (drop, spray and biocontrol+ spray) at different time intervals (24, 48, 72, 96 and 120 h) after pathogen inoculation.




Fig 5. Development of initial black/brown disease lesions on the detached leaves of *B. juncea* variety RLC-3 using different pathogen inoculation methods (drop, spray and biocontrol+ spray) at different time intervals (24, 48, 72, 96 and 120 h) after pathogen inoculation.

respectively after 120 hrs (Fig. 2). The drop method showed the highest number of initial lesions followed by spray (Fig. 3).

The *B. juncea* (2nd variety- RLC-3) at 24 h after inoculation of pathogen, drop method showed highest number of initial lesions that is (14) lesions, followed by spray (5) lesions and biocontrol+spray (0). Maximum number of lesions i.e. 59, 47 and 1 were appear on leaves of *B. juncea* using drop, spray and biocontrol+spray method, respectively after 120 hrs (Fig. 4). The drop method showed the highest number of initial lesions followed by spray (Fig. 5).

In *B. napus* (Variety- 'GSC-7') at 24 h after inoculation of pathogen, drop method showed highest number of initial lesions that is (10) lesions, followed by spray (6) and biocontrol+ spray (0). Maximum number of lesions i.e. 53, 35 and 2 were appear on leaves of *B. napus* using drop, spray and biocontrol+spray method, respectively after 120 hrs (Fig. 6). The drop method showed the highest number of initial lesions followed by spray (Fig. 7).

The *B. nigra* (Variety- 'PC-6') at 72 h after inoculation of pathogen, drop method showed highest number of initial lesions that is (44) lesions, followed by spray (19) lesions and biocontrol+spray (3). Maximum number of lesions i.e. 73, 55 and 4 were appear on leaves of *B. nigra* using drop, spray and biocontrol+spray method, respectively after 120hrs (Fig. 8). At 48, 72, 96, and 120 h after inoculation of pathogen, thereby the drop method showed the highest number of initial lesions followed by spray (Fig. 9).

Results revealed that out of three inoculation methods used, spore suspension drop inoculation method was most ideal as this fixed the inoculum on the target site. With this technique, a single conidiophore of *A. brassicae* was able to cause local lesions on leaves. This was also reported by Xu and Ko (1998), that even a single conidium containing drop were able to induce the development of local lesion in black mustard leaves. It also produces significantly higher number of disease lesions than the other inoculation methods and has the advantage of being accurate and precise. It is also easy to handle the inoculated plants. The results indicated that two varieties *B. juncea* (RLC-3) and *B. napus* (GSC-7) showed some tolerance as compared to varieties *B. juncea* (Giriraj) and *B. nigra* (PC-6). The biocontrol agent used in particular study showed that the tolerance can be induced in the plants using *T. viride*. As the number of lesions and pathogenicity caused by *A. brassicae* was reduced considerably when the culture of *T. viride* was used.




Fig. 6. Number of lesions on the detached leaves of *B. napus* (variety-GSC-7) using different pathogen inoculation methods (drop, spray and biocontrol+ spray) at different time intervals (24, 48, 72, 96 and 120 h) after pathogen inoculation.




Fig. 7. Development of initial black/brown disease lesions on the detached leaves of *B. napus* variety GSC-7 using different pathogen inoculation methods (drop, spray and biocontrol+ spray) at different time intervals (24, 48, 72, 96 and 120 h) after pathogen inoculation.

References

- Allen SJ, Brown JF and Kochman JK. 1983. Effect of leaf age, host growth stage, leaf injury, and pollen on the infection of sunflower by *A. helianthi*. *Phytopathol* **73**: 896-898.
- Anonymous 2016. https://nmoop.gov.in/Publication/StatusPaper_RandM_2017.pdf.




Fig. 8. Number of lesions on the detached leaves of *B. nigra* (variety-PC-6) using different pathogen inoculation methods (drop, spray and biocontrol+ spray) at different time intervals (24, 48, 72, 96 and 120 h) after pathogen inoculation.




Fig 9. Development of initial black/brown disease lesions on the detached leaves of *B. nigra* variety PC-6 using different pathogen inoculation methods (drop, spray and biocontrol+ spray) at different time intervals (24, 48, 72, 96 and 120 h) after pathogen inoculation.

Buckley PM, Williams WP and Windham GL. 2009. Comparison of two inoculation methods for evaluating corn for resistance to aflatoxin contamination. Mississippi Agricultural & Forestry Experiment Station, Mississippi State University, Bulletin 1148.

Co O, Ngugi K, Nzioki H and Githiri SM. 2008. Evaluation of smut inoculation techniques in sugarcane seedlings. *Proc S Afr Sug Technol Assoc* **81**:478-481.

Elad Y. 1994. Biological control of grape graymold by *Trichoderma harzianum*. *Crop Prot* **13**: 35-38.

Giri P, Taj G and Kumar A 2013. Comparison of artificial inoculation methods for pathogenesis studies of *A. brassicae* (Berk.) Sacc. on Indian mustard [*B. juncea* (L.) Czern. & Coss.]. *Afr J Biotechnol* **12**: 2422-2426.

Goyal P, Chahar M, Mathur AP, Kumar A and Chattopadhyay C. 2011. Morphological and cultural variation in different oilseed Brassica isolates of *Alternaria brassicae* from different geographical regions of India. *Indian J Agric Sci* **81**: 1052-1058.

Hong CX and Fitt BDL. 1995. Effect of inoculum concentration, leaf age and wetness period on the development of dark leaf and pod spot (*Alternaria brassicae*) on oilseed rape (*B. napus*). *Ann Appl Biol* **127**: 283-295.

Jie L, Zifeng W, Lixiang C, Hongming T, Patrik I, Zide J and Shining Z. 2009. Artificial inoculation of banana tissue culture plantlets with inigenous endophytes originally derived from native banana plants. *Biol Control* **51**:427-434.

Khan MM, Khan RU and Mohiddin FA. 2007. Variation in occurrence and morphology of *A. brassicae* (Brek) Sacc. Causing blight in rapeseed and mustard. *Ann Pl Protec Sci* **15**: 414-417.

Kumar S, Upadhyay JP and Kumar S. 2005. Biocontrol of *Alternaria* leaf spot of *Vicia faba* using antagonistic fungi. *J Biol Control* **20**: 247-251.

Lapwood DH and Mckee RK. 1966. Dose - response relationships for infection of potato leaves by zoospores of *Phytophthora infestans*. *Trans Br Mycol Soc* **49**: 679-686.

Louvet J.1958. The black spot disease of colza *A. brassicae*. *CR Acad Agric Fr* **44**: 694.

McDonald WC.1959. Gray leaf spot of rape in Manitoba. *Can J Plant Sci* **39**: 409.

Meena PD, Awasthi RP, Chattopadhyay C, Kolte SJ and Kumar A. 2010. *Alternaria* blight: a chronic disease in rapeseed-mustard. *J Oilseed Brassica* **1**: 1-11.

Meena PD, Rani, A, Meena R, Sharma P, Gupta R and Chowappa P. 2012. Aggressiveness, diversity and distribution of *A. brassicae* isolates infecting oilseed Brassica in India. *Afr J Microbiol Res* **6**: 5249-5258.

Meena PD, Meena RL, Chattopadhyay C and Kumar A.2004. Identification of critical stage for disease development and biocontrol of *Alternaria* blight of Indian mustard (*B. juncea*). *J Phytopathol* **152**: 204-209.

- Mohiddin FA, Khan MR, Khan SM and Bhat BH. 2010. Why *Trichoderma* is considered super hero (super fungus) against the evil parasites? *Plant Path J* **9**: 138-148.
- Perello AE, Moreno MV, Monaco C, Simon MR and C Cordo. 2009. Biological control of *Septoria tritici* blotch on wheat by *Trichoderma spp.* under field condition in Argentina. *Biol Control* **54**: 113-122.
- Prasad L and Vishnuvat K. 2006. Assessment of yield loss in cauliflower seed crop due to *Alternaria* blight. *Indian Phytopath* **59**: 185-189.
- Reshu and Khan MM. 2012. Role of different microbial origin bioactive antifungal compounds against *Alternaria spp.* causing leaf blight of mustard. *Pl Pathol J* **11**: 1-9.
- Reshu and Khan MM. 2012. Role of different microbial origin bioactive antifungal compounds against *Alternaria spp.* causing leaf blight of mustard. *Pl Pathol J* **11**: 1-9.
- Shah FA, Wang CS and Butt TM. 2005. Nutrition influences growth and virulence of the insect pathogenic fungus *Metarhiziuma nisopliae*. *FEMS Microbiol Lett* **251**: 259-266.
- Shah SA, Iftikhar A, Rahmkan K and Mumtaz A. 2005. 'NIFA-mustard canola' first mutant variety of oilseed mustard [*B. juncea* (L.) Czern & Coss.] in Pakistan. *Mutat Breed-Newslett Rev* **1**: 22-23.
- Singh B, Kaur T, Kaur S and Kaur A. 2013. Insecticidal potential of an endophytic *Cladosporium velox* against *Spodopteralitura* mediated through inhibition of alpha glycosidase. *Pest Biochem Physiol* DOI: 10.1016/j.pestbp.2016.01.004.
- Singh B, Thakur A, Kaur S, Chadha BS and Kaur A. 2012. *Acetylcholinesterase* inhibitory potential and insecticidal activity of an endophytic *Alternaria sp.* from *Ricinus communis*. *Appl Biochem Biotechnol* **168**: 991-1002.
- Toussoun TA and Nelson PE 1976. A pictorial guide to the identification of *Fusarium Species*, second edition. Pennsylvania State University Press, University Park. 43 pp.
- Toussoun TA, Nash SM and WC Snyder. 1960. The effect of nitrogen sources and glucose on the pathogenesis of *Fusarium solani f. phaseoli*. *Phytopathol* **50**: 137-140.
- Tuite J. 1969. Plant Pathological Methods - Fungi and Bacteria. *Burgess Publ Co., Minneapolis, Minnesota*, 239 pp.
- Xu XL and Ko WH. 1998. A quantitative confined inoculation method for studies of pathogenicity of fungi on plants. *Bot Bull Acad Sin* **39**: 187-190.