



## Response of bioregulators and irrigation on plant height of Indian mustard (*Brassica juncea* L.)

Ajeet Singh\* and Ram Swaroop Meena

Department of Agronomy, Institute of Agricultural Sciences, Banaras Hindu University,  
Varanasi-221005 Uttar Pradesh, India

\*Corresponding author: [ajitbhu89@gmail.com](mailto:ajitbhu89@gmail.com)

(Received: 15 November 2019; Revised: 12 December 2019; Accepted: 16 December 2019)

### Abstract

The field experiment was carried out during the 2014-15 and 2015-16 in *Rabi* season to evaluate the response of bioregulators and irrigation on growth parameters of Indian mustard (*Brassica juncea* L.). The treatments consisted three irrigation levels (no irrigation, one irrigation and two irrigation) and seven foliar spray of bioregulators (control (water spray), thiourea @ 1000ppm, thiourea @ 500ppm, salicylic acid @ 100ppm, salicylic acid @ 50ppm, glycinebetaine @ 100ppm and glycinebetaine @ 50ppm). The result indicated that in the irrigated plots significantly improved the plant height of Indian mustard. The application of two irrigation recorded highest plant height at 80 DAS (132.7, 119.9 and 126.3 cm), 100DAS (173.6, 168.5 and 171.1 cm) and at harvest (176.4, 171.9 and 174.2 cm) followed by the application of one irrigation in both years and pooled analysis, respectively. Moreover, the application of bioregulators significantly improves the plant height of the Indian mustard. The application of thiourea @ 1000ppm gave highest plant height at 80 DAS (130.5, 118.2 and 124.4 cm), 100DAS (172.4, 166.8 and 169.6 cm) and at harvest (175.3, 170.9 and 173.1 cm). However, the application of thiourea @ 500 ppm was statistically at par with thiourea @ 1000ppm in both years and pooled analysis, respectively. Furthermore, combined result of bioregulators and irrigation levels were observed significant on plant height of Indian mustard. The application of two irrigation+ foliar spray of thiourea @ 1000ppm recorded the highest plant height at 100DAS (190.2, 182.3 and 186.3 cm) and at the time of harvest (194.2, 187.6 and 190.9 cm) over control. The application of two irrigation+ foliar spray of thiourea @ 500ppm was statistically at par at 100 DAS (189.4, 181.6 and 185.5 cm) and at harvest (193.3, 186.7 and 190.0 cm) with two irrigation+ foliar spray of thiourea @ 1000ppm in both years and pooled analysis, respectively. The present study highlighted the practical importance of the irrigation levels and foliar spray of bioregulators combination response on plant height of Indian mustard.

**Key words:** Bioregulators, growth, Indian mustard, irrigation, plant height, thiourea

### Introduction

India is one of the leading oilseeds producing country in the globe. Oilseeds crop are the succeeding largest agricultural commodity after cereals. Indian mustard [*Brassica juncea* (L.) Czern & Coss.] is the second important edible oilseed crop after groundnut, meeting the fat requirement of about 50 per cent population in all the northern states (Shivran *et al.*, 2019). The mustard oil cake contains 5.1 per cent nitrogen, 1.8 per cent phosphorus and 1.2 percent potassium and it rich sources of protein (40 per cent) and it grown in area of about 68.57 lakh ha with production of 69.74 lakh tones (DES, 2018). Due to scarcity of winter rainfall mustard shows better response to irrigation. This crop is more sensitive to water fluctuation and more or less at critical growth stages, which adversely influenced the yield (Meena *et al.*, 2013). The irrigation at critical stages increased the yield of the Indian mustard this might due to the more availability of the nutrient and more efficient metabolic activities of the

plant (Mehta, 2004). Moreover, it is universally accepted that limited irrigation water can be best be utilized more efficiently by scheduling irrigation at critical growth stages of the crop (Saud *et al.*, 2016).

Rapeseed and mustard is generally, affect by irrigation water. The rapeseed-mustard is the crop that has fewer requirements of the water (Aujla *et al.*, 2005, Meena *et al.* 2017). To alleviate the harmful effect of drought stress, bioregulators may be used to change the different metabolic and physiological activities of the plant for increasing the yield of mustard crop (Jat, 2007). Thiourea containing one SH group and it's is a sulphhydryl compound. The SH group has been implicated in photosynthetic translocation in plants (Giaquinta, 1976). It is simply concerned in light activation of photosynthetic enzymes (Salisbury and Ross, 1986). The exogenous application of Salicylic acid (SA) has been reduced the negative effect of water stress (Khan *et al.*, 2015, Meena *et al.*, 2018) and spray of SA improve the growth of the

plant (Hayat *et al.*, 2010). Glycinebetaine an important ammonium compound, it considered to be one of the most pre-dominant and effective osmoprotectants. The exogenous application of glycinebetaine improved the drought tolerance in plants (Mahmood *et al.*, 2009). Hence, the study was conducted to see the effect of bioregulators on dry matter accumulation of Indian mustard under different levels of irrigation.

## Materials and Methods

An experiment was conducted during the *rabi* (winter) seasons of 2014-15 and 2015-16 at the Agricultural Research Farm, Banaras Hindu University, Varanasi. The experimental field has clay loam soil. The ploughing layer of soil has pH (7.58 and 7.60) organic carbon (0.44% and 0.46%), available nitrogen (217.3 and 224.1 kg ha<sup>-1</sup>), available phosphorus (20.28 and 22.85 kg ha<sup>-1</sup>), available potassium (219.2 and 226.4 kg ha<sup>-1</sup>) and available sulphur (20.90 and 21.85 mg kg<sup>-1</sup> ha<sup>-1</sup>) during both the years, respectively. The field experiment was laid out in split plot design. The experiment have 21 treatments combinations *viz.*, Main plot treatment– three irrigation levels (zero irrigation, one irrigation and two irrigation) and Sub-plots treatment– seven bioregulators spray (control, thiourea @1000ppm, thiourea @ 500ppm, salicylic acid @ 100ppm, salicylic acid @ 50ppm, glycinebetaine @ 100ppm and glycinebetaine @ 50ppm) and irrigation will apply at pre-flowering and pod formation stage @ 6 cm by V-notch method and Foliar spray done at 50 per cent flower initiation and 50 per cent pod formation stage. Furrows were opened at a spacing of 45×15 cm for the sowing of Indian mustard variety ‘Ashirwad’ with seed rate of 5 kg ha<sup>-1</sup>. NPK and S were applied as 100, 50, 50 and kg ha<sup>-1</sup> by basal application. Nitrogen and phosphorus were applied through diammonium phosphate and urea, potash and sulphur through murate of potash and elemental sulphur, respectively. The data were analyzed as per the standard procedure for “Analysis of Variance” (ANOVA) (Gomez and Gomez, 1976).

## Results and Discussion

### Response of irrigation

The revealed of the data present in table 1 indicated that the application of irrigation levels significantly increased the plant height of the Indian mustard over control. The maximum plant height (132.7, 119.9 and 126.3 cm at 80 DAS; 173.6, 168.5 and 171.1 cm at 100DAS cm and 176.4, 171.9 and 174.2 cm at harvest) was recorded with the application of two irrigation (at pre flowering and pod formation stage) in both the years and pooled analysis. This might be due to the adequate soil moisture increase

the availability of the nutrient in the soil for the plant to increase in growth parameters by cell elongation and cell division this ascribed due to higher activity of auxin in plant tissues and photosynthesis activity of plant so they produced more food (Mandal *et al.*, 2006). The similar result recognized by Lal *et al.* (2000), Sharma *et al.* (2006), Nagdive *et al.* (2007), Mandal *et al.* (2010) and Meena *et al.* (2018a).

### Response of bioregulators

The data present in table 1 showed that the application of bioregulators significantly increased the plant height of the Indian mustard. The higher value of plant height at 80 DAS (130.5, 118.2 and 124.4 cm), 100DAS (172.440, 166.8 and 169.6 cm) and at harvest (175.3, 170.9 and 173.1 cm) was recorded with the thiourea @1000ppm and it's found statically at par with the application of thiourea @500ppm in both the years and pooled analysis. This may be ascribed due to the foliar application of thiourea motivating the photosynthetic carbon fixation mechanism that enhanced the canopy of the plant (Mehta and Sumeria, 2001; Meena *et al.*, 2019). Thiourea exhibits cytokinin activity (Deshveer and Singh, 2003) and cytokinin delaying the leaf senescence of the plant (Solanki, 2003). The similar result reported by Sahu *et al.* (2005), Srivastava *et al.* (2010), Singh *et al.* (2017) and Meena *et al.* (2020).

### Interaction

The examination of data in table 2 and 3 indicated that the Interaction effect of levels of irrigation and foliar spray of bioregulators were observed significant on plant height of Indian mustard. Result showed that two irrigation was recorded maximum plant height at 100 days after sowing (190.2, 182.3 and 186.3 cm) and at the time of harvest (194.2, 187.6 and 190.9 cm) with the application of thiourea @1000ppm and its was statistically at par with two irrigation with thiourea @ 500ppm at 100 DAS (189.4, 181.6 and 185.5 cm) and at harvest (193.3, 186.7 and 190.0 cm) during both the years and pooled analysis. While, the minimum plant height at 100 DAS (84.8, 81.2 and 82.9 cm) and at harvest (85.8, 82.0 and 83.9 cm) was recorded in zero irrigation + water spray during both the years and pooled analysis, respectively. The similar result reported by Muhl *et al.*, (2014), Muhl and Solanki (2015), Hassanein *et al.* (2012), Meena (2020) and Meena (2020a).

## References

- Aujla MS, Thind HS and Buttar GS. 2005. Cotton yield and water use efficiency at various levels of water and N through drip irrigation under two methods of planting. *Agril Water Manag* **71**:167-179.
- Chimenti CA, Pearson J and Hall AJ. 2002. Osmotic adjustment and yield maintenance under drought in sunflower. *Field Crop Res* **10**:235-246.

Table 1. Effect of irrigation levels and foliar spray of bio-regulators on plant height of Indian mustard

Treatment	Plant height (cm)														
	40 DAS			60 DAS			80 DAS			100 DAS			harvest		
	2014-15	2015-16	Pooled	2014-15	2015-16	Pooled	2014-15	2015-16	Pooled	2014-15	2015-16	Pooled			
<b>Irrigation levels</b>															
Zero irrigation	53.5	50.6	52.1	78.1	73.0	75.5	103.2	93.0	98.1	129.3	128.7	129.0	131.5	131.1	131.3
One irrigation	53.7	50.8	52.2	78.3	73.2	75.8	126.6	113.9	120.3	163.8	160.5	162.1	166.1	162.9	164.5
Two irrigation	55.0	52.0	53.5	80.2	75.0	77.6	132.7	120.0	126.3	173.6	168.5	171.1	176.4	171.9	174.2
SE $\pm$	1.1	1.0	0.7	1.5	1.4	1.1	2.6	2.9	1.9	1.8	1.6	1.2	2.0	1.8	1.3
CD ( $p=0.05$ )	NS	NS	NS	NS	NS	NS	10.1	11.4	6.3	7.1	6.3	3.9	7.8	7.0	4.4
<b>Foliar spray of bio-regulators</b>															
Control (water spray)	53.4	50.5	52.0	77.9	72.9	75.4	98.7	89.0	93.9	118.3	119.5	118.9	120.6	119.6	120.1
Thiourea @ 500 ppm	54.6	51.6	53.1	79.6	74.4	77.0	128.3	117.8	123.0	171.3	165.8	168.6	174.1	169.8	171.9
Thiourea @ 1000 ppm	54.7	51.7	53.2	79.8	74.6	77.2	130.5	118.2	124.4	172.4	166.8	169.6	175.3	170.9	173.1
Salicylic acid @ 50 ppm	53.9	51.0	52.5	78.7	73.6	76.1	121.5	109.7	115.6	159.9	156.4	158.2	162.1	159.4	160.7
Salicylic acid @ 100 ppm	54.4	51.4	52.9	79.4	74.2	76.8	127.9	115.9	121.9	165.4	161.2	163.3	168.3	164.9	166.6
Glycinebetaine @ 50 ppm	53.4	50.5	52.0	78.0	72.9	75.4	110.7	99.8	105.3	146.1	145.0	145.6	147.8	147.0	147.4
Glycinebetaine @ 100 ppm	54.0	51.0	52.5	78.8	73.6	76.2	116.4	104.4	110.4	153.5	151.1	152.3	155.6	153.8	154.7
SE $\pm$	1.4	1.3	0.9	2.0	1.9	1.4	1.6	1.7	1.2	0.9	0.8	0.6	1.1	0.9	0.7
CD ( $p=0.05$ )	NS	NS	NS	NS	NS	NS	4.6	4.7	3.2	2.7	2.3	1.7	3.2	2.6	2.0

Table 2. Interactive effect of foliar spray of bioregulators and irrigation levels on plant height (cm) of Indian mustard at 100DAS

Treatment	2014-15				2015-16				pooled
	No irrigation	One irrigation	Two irrigation	No irrigation	One irrigation	Two irrigation	No irrigation	One irrigation	
Control (water spray)	84.8	128.3	141.9	81.2	135.3	142.2	83.0	131.8	142.1
Thiourea @ 500 ppm	147.6	179.9	189.4	145.7	173.2	181.6	146.6	176.6	185.5
Thiourea @ 1000 ppm	149.1	180.9	190.2	146.9	174.1	182.3	148.0	177.5	186.3
Salicylic acid @ 50 ppm	135.2	167.4	177.3	134.5	163.0	171.7	134.8	165.2	174.5
Salicylic acid @ 100 ppm	140.2	173.3	182.7	138.9	168.2	176.5	139.6	170.8	179.6
Glycinebetaine @ 50 ppm	120.7	155.1	162.5	124.4	152.1	158.6	122.5	153.6	160.6
Glycinebetaine @ 100 ppm	127.6	161.4	171.3	129.0	157.7	166.5	128.3	159.6	168.9
SE <sub>me</sub> ±	1.6			1.4			1.1		
CD ( <i>p</i> =0.05)	4.6			3.96			2.98		

Table 3. Interactive effect of foliar spray of bioregulators and irrigation levels on plant height (cm) of Indian mustard at harvest

Treatment	2014-15				2015-16				Pooled
	No irrigation	One irrigation	Two irrigation	No irrigation	One irrigation	Two irrigation	No irrigation	One irrigation	
Control (water spray)	85.8	133.6	142.5	82.0	134.5	142.3	83.9	134.0	142.4
Thiourea @ 500 ppm	149.5	182.4	193.3	148.4	177.2	186.7	149.0	179.8	190.0
Thiourea @ 1000 ppm	151.2	183.5	194.2	149.9	178.2	187.5	150.5	180.9	190.9
Salicylic acid @ 50 ppm	136.4	169.3	180.5	137.0	165.7	175.6	136.7	167.5	178.1
Salicylic acid @ 100 ppm	142.2	176.0	186.7	142.0	171.6	181.0	142.1	173.8	183.9
Glycinebetaine @ 50 ppm	124.7	155.2	163.7	126.7	153.4	160.8	125.7	154.3	162.2
Glycinebetaine @ 100 ppm	130.7	162.4	173.8	131.9	159.7	169.6	131.3	161.1	171.7
SE <sub>me</sub> ±	1.96			1.6			1.3		
CD ( <i>p</i> =0.05)	5.6			4.5			3.5		

- DES. 2017. Directorate of Economics and Statistics, Ministry of Agriculture, www.agri.com
- Deshveer CL and Singh A. 2003. Effectiveness of agrochemicals Indian mustard (*B. juncea*) *Indian J Agril Sci* **73**:345-346.
- Gaiser T, Barros DI, Lange FM and Williams JR. 2004. Water use efficiency of a maize/cowpea intercrop on a highly acidic tropical soil as affected by liming and fertilizer application. *Plant and Soil* **263**:165-171.
- Giaquinta RT. 1976. Evidence of phloem loading from the apoplast: Chemical modification of membrane sulphhydryl group. *Plant Physiol* **58**:872-875.
- Gomez KA and Gomez AA. 1976. Statistical Procedures for Agricultural Research, 2nd edn. John Willey and Sons Inc. New York, USA.
- Hassanein RA, Abdelkader AF, Ali H, Amin AAE and Rashad E.M. 2012. Grain primary and foliar pretreatment enhanced stress defense in wheat (*Triticum aestivum* var. Gimaza) plants cultivated in drought land. *Aus J Crop Sci* **6**: 121-129.
- Hayat Q, Hayat S, Ifran M and Ahmad A. 2010. Effect of exogenous salicylic acid under changing environment: a review. *Envl Expl Bot* **68**:14-25.
- Iqbal N and Ashraf M. 2009. Glycine betaine, an osmolyte of interest to improve water stress tolerance in sunflower (*Helianthus annuus* L.): water relation and yield. *South African J Bot* **74**:274-281.
- Jat RP. 2007. Effect of organic and inorganic sources of nutrients, gypsum and thiourea on growth, yield and quality of mustard. M.Sc. (Ag) Thesis, Department of Agronomy, MPUAT, Udaipur.
- Khan MIR, Fatma M, Per TS, Anjum NA and Khan NA. 2015. Salicylic acid on photosynthetic pigments and antioxidative enzymes of soybean plant. *Front Pl Sci* **6**:1-17.
- Lal R, Singh S, Yadav SK, Yadav DB and Chand M. 2000. Influence of irrigation and nitrogen levels on the quality and oil yield of Brassica species. *Ann Plant Soil Res* **2**:85-88.
- Mahmood T, Asraf M and Shahbaz M. 2009. Does exogenous application of glycine betaine as a pre-sowing seed treatment improve growth and regulate some key physiological attributes in wheat plants grown under water deficit condition. *Pak J Bot* **41**:1291-1302.
- Mandal KG, Hati KM, Misra AK and Bandyopadhyay KK. 2010. Root biomass, crop response and water-yield relationship of mustard (*B. juncea* L.) grown under combinations of irrigation and nutrient application. *Irrigat Sci* **28**:271-280.
- Mandal KG, Hati KM, Misra AK and Bandyopadhyay KK. 2006. Assessment of irrigation and nutrient effects on growth, yield and water use efficiency of Indian mustard (*B. juncea*) in central India. *Agril Water Manag* **85**:279-286.
- Meena RK, Sharma DD, Chouhan GS, Shukla KB, Sumeriya HK and Padiwal NK. 2013. Growth and productivity of mustard [*B. juncea* (L.) Czern & Coss.] as influenced by irrigation levels and agrochemicals. *Ann Agri-Bio Res* **18**:312-317.
- Meena RS, Kumar S and Yadav GS. 2020. Soil carbon sequestration in crop production. In: Meena RS (Eds) Nutrient dynamics for sustainable crop production. Springer, Singapore. P: 1-39.
- Meena RS, Kumar S, Bohra JS, Lal R, Singh GS and Pandey A. 2019. Response of alley cropping grown sesame to lime and sulphur on yield and available nutrient status in an acidic soil of Eastern India. *Energy Ecol Env* **4**: 65-74.
- Meena RS, Kumar V, Yadav GS and Mitran T. 2018. Response and interaction of *Bradyrhizobium japonicum* and *Arbuscular mycorrhizal* fungi in the soybean rhizosphere: A review. *Pl Growth Regulat* **84**:207-223.
- Meena RS, Meena PD, Yadav GS and Yadav SS. 2017. Phosphate solubilizing microorganisms, principles and application of microphos technology. *J Cleaner Product* **145**: 157-158.
- Meena RS, Mitran T, Kumar S, Yadav G, Bohra JS and Datta R. 2018a. Application of remote sensing for sustainable agriculture and forest management. *Informat Process Agri* **5**: 295-297.
- Meena RS. 2020. Nutrient Dynamics for Sustainable Crop Production. Springer, Singapore. ISBN: 978-981-13-8659-6 DOI: <https://doi.org/10.1007/978-981-13-8660-2>
- Meena RS. 2020a. Soil Health Restoration and Management. Springer, Singapore. ISBN: 978-981-13-8569-8 DOI: <https://doi.org/10.1007/978-981-13-8570-4>
- Mehta JP. 2004. Response of bioregulators in mitigating moisture stress in mustard [*B. juncea* (L.) Czern & Coss]. Ph.D Thesis, Department of Agronomy, RCA, MPUAT, Udaipur.
- Mehta YK and Sumeria HK. 2001. Effect of plant growth regulators sulphuric acid on dry matter partitioning of pea cultivars. *Haryana J Agron* **17**:173-174.
- Muhal S and Solanki NS. 2015. Effect of seeding dates and salicylic acid foliar spray on growth, yield, phenology and agro-meteorological indices of Brassica species. *J Oilseed Brassica* **6**:183-190.

- Muhal S, Solanki NS, Singh P and Shukla KB. 2014. Effect of salicylic acid on productivity and nutrient uptake of Brassica species under different planting durations. *African J Agril Res* **9**:1101-1106.
- Nagdive SJ, Bhalerao PD, Dongarwar UR and Gond VV. 2007. Effect of irrigation and nutrient management on yield, quality and uptake of nutrients by mustard (*B. juncea* L.). *J Soil and Crops* **17**:128-132.
- Sahu MP, Kumawat SM, D'Souza SF, Ramaswamy NK and Singh G. 2005. Sulphydryl bioregulator technology for increasing mustard production. *Res Bulletin RAU-BARC* 1-52.
- Salisbury FB and Ross CW. 1986. *Plant Physiology*. Wads Worth Publishing Company, USA, 206-207.
- Saud RK, Singh BP and Pannu RK. 2016. Studied on consumptive use, water use efficiency and moisture extraction pattern by Indian mustard as influenced by limited irrigation and nitrogen levels. *Intl J For Crop Improv* **7**:137-140.
- Sharma SK, Mendhe SN, Kolte HS, Rajput GR and Yenpreddiwar MD. 2006. Effect of sowing and irrigation management on growth and yield of mustard (*B. juncea* L.). *J Soil Crops* **16**:455-459.
- Shivran A, Patel BJ and Gora M. 2019. Effect of irrigation schedule and bio-regulators on yield attributes and yield of mustard [*B. juncea* (L.) Czern & Coss] crop. *Intl J Chem Studies* **7**:1874-1877.
- Singh A, Singh AK and Ashwin C. 2017. Effect of hydrogel and thiourea on yield, quality and nutrient uptake of Indian mustard under moisture stress condition. *Res on Crops* **18**:42-48.
- Solanki NS. 2003. Effect of thiourea and dimethyl sulphoxide on phosphorus use efficiency, dry matter partitioning and productivity of clusterbean [*Cyamopsis tetragonolob* (L.) Taub.]. Ph.D. Thesis, MPUAT, Udaipur, Rajasthan.
- Srivastava AK, Ramaswamy NK, Suprasanna P and D'Souza SF. 2010. Genomewide analysis of thiourea modulated salinity stress responsive transcripts in seeds of *B. juncea*. Identification of signalling and effector components of stress tolerance. *Ann Bot (London)* **106**: 663-74.