



Biochemical characterization of Indian mustard (*Brassica juncea* L.) genotypes in response to moisture stress and irrigation modules

Sukhmaninder Kaur¹ and Pushp Sharma^{2*}

¹Department of Botany ²Department of Plant Breeding and Genetics, Punjab Agricultural University, Ludhiana, Punjab 141004 India

*Corresponding author: pushp20@yahoo.com; pushp20@pau.edu

(Received: 15 May 2015; Revised: 15 June 2015; Accepted: 26 June 2015)

Abstract

In the current investigation, the impact of moisture stress and irrigation modules was studied on biochemical characteristics in twelve *Brassica juncea* genotypes. Experiment was laid in split plot design with three moisture treatments viz. only pre sowing irrigation (I₀), one irrigation at 35 DAS (I₁) and two irrigations, one at 35 DAS and second at 65 DAS (I₂). Water stress up-regulated the sugars and proline content in the genotypes while it reduced the protein content. Highest amount of total sugars was recorded in K-9-108 (70.3 mg g⁻¹ DW), reducing sugars in RLC1 (12.61 mg g⁻¹ DW), protein content in MLM-19 and NLM-3 (8.8 mg g⁻¹ DW) while highest proline was in NPJ-79 (0.87 mg g⁻¹ DW) under moisture stress. Genotypes differed significantly for the biochemical constituents and total soluble proteins increased progressively with one and two irrigations.

Key words: Irrigation modules, moisture stress, proline, reducing sugars, sugars

Introduction

Water stress is one of the important and limiting factor that affects plant growth and productivity in the arid and semi-arid regions of the world. Drought causes a severe impairment in plant photosynthesis, growth and development and hence limits plant production and performance of crop plants worldwide (Azadeh *et al.*, 2014). Water deficit leads to reduced nutrient uptake by roots and transportation from roots to shoots, due to restricted transpiration and impaired active transport and membrane permeability (Yuncaï and Schmidhalter, 2005). Also, the response of plants to water stress differ significantly depending upon intensity and duration of the stress, plant species and plant growth stage (Jaleel *et al.*, 2008). In response to water deficit, plants evolve biochemical adaptations and exhibit several alterations in metabolic processes viz. accumulation of low molecular weight sugars, amino acids or betaines which maintain cellular turgor as a consequence of decreased water potential. Therefore, at the cellular level, plants attempt to alleviate the damaging effects of stress by

altering their metabolism to cope with stress (Bayoumi *et al.*, 2008). The decreased water availability negatively affects the metabolite concentration, followed by alteration in carbohydrate metabolism and increased synthesis of compatible solutes such as reducing sugars. The organic and inorganic solutes thus accumulated raises the osmotic pressure in the cytosol, thereby maintaining cellular turgor and a driving force for water uptake. The level of sugars generally increases under water for removal of closely associated water from the protein without leading to conformational changes and loss of enzymatic functions (Yordonov *et al.*, 2003). An increase in sucrose and hexose levels under moisture stress has been proposed as the osmotic sugars, adjustment in sucrose transporting species (Westgate and Boyer, 1985). On the other hand decrease in protein synthesis causes rapid dehydration. Therefore, changes in protein content is considered as an important response of plants to environmental stress and as an adaptive response towards moisture stress. Earlier reports the positive (Shahraki *et al.*, 2008, Tohidi *et al.*, 2011) as well

as non-significant effect of drought (Praveen *et al.*, 1996, Tahir *et al.*, 2007) on protein content which are affected by various factors like variety, class and environmental stress encountered during plant growth and development. Ahmadi *et al.* (2010) reported increased protein content in maize seedlings exposed to mild water stress which decreased on exposure to severe drought. Drought tolerance is an interactive association of complex morphological, physiological and molecular characters associated with low molecular weight biomolecules like proline, the most compatible osmolyte increasing under drought stress and is considered as an important stress tolerance mechanism (Verbruggen and Hermans, 2008). Biochemical alterations and the adaptive response of plants towards water stress tolerance is of great importance for the plant breeders in developing drought tolerant varieties by understanding the detrimental effect of drought on biochemical traits in plants. The present study was therefore undertaken to visualize the alterations in the biochemical characters in Indian mustard genotypes as influenced by moisture deficit and irrigation modules.

Material and Methods

Thirty genotypes of *B. juncea* were evaluated under field conditions of oilseeds farms (30° 54'N 75°48'E 247m) located at Punjab Agricultural University Ludhiana, Punjab, India. Based on their performance twelve genotypes were selected for further study during two consecutive years 2009-10 and 2010-11. Experiments were laid down in split plot design in three replications according to recommendations of package of practices. Treatments comprised of irrigation schedule in main plots and genotypes in subplots. Irrigation regimes consisted of only pre-sowing irrigation designated as moisture stress (I_0), one irrigation at 35DAS (I_1 , restricted moisture) and two irrigations at 35 and then at 65DAS (I_2 , normal moisture). Three plants were tagged per treatment in each replication. 3rd and 4th leaf of the main shoot from the tagged plants were collected at 90 DAS. The sampled leaves were oven dried at 60 °C ± 1°C for 24 hrs and were used for biochemical estimations. Standard protocols were followed to estimate total

soluble sugars (Dubois *et al.*, 1956), reducing sugars (Nelson, 1944), total soluble proteins (Lowry *et al.*, 1951) and Proline (Bates *et al.*, 1973). The difference between total soluble sugars and reducing sugars was computed which gave the non-reducing sugar content in the cultivars. The CPCS1 software developed at PAU was applied for statistical analysis. The effects were computed at 5% and 1% level of significance.

Results and Discussion

Variations existed between the genotypes studied. The irrigation modules and the interactive effects (genotypes x irrigations) were significant for all the sugar moieties.

Sugars: Total sugar content was highest in cultivar K-9-108 under stress and in cultivar QM-7-196 both under restricted and normal moisture regimes. K-109-113 possessed lowest total sugar content under all moisture regimes (Fig.1). Total soluble sugars acts as osmoprotectant thus maintaining the turgor pressure and stabilizes the cellular membranes. Soluble carbohydrates play a potential role in adaptation to water stress as reported in maize (Mohammadkhani and Heidari, 2008). Increased sugar content under drought stress has been reported in maize by Sinay and Karuwal (2014) and Homayouni and Khazarian (2014) as well in *Brassica napus* (Nosrati *et al.*, 2014). RLC1 had highest reducing sugars under all the moisture regimes. Maximum amount of non-reducing sugars were estimated in K-9-108 (62.6 mg g⁻¹ DW) under moisture stress while QM-7-196 registered highest content under restricted and normal moisture schedules. K-9-108 had the lowest content of non-reducing sugars under 3 irrigation regimes (Fig.2). Overall, the content of total sugars, reducing and non-reducing sugars was highest cultivars under moisture stress and the decline in respective content was observed with one and two irrigation module.

Total soluble proteins (TSP): Genotypes differed significantly for the total soluble protein content but their content increased with irrigations applied. Cultivars registered maximum soluble proteins with two sequential irrigations given at 35 and 65 DAS.




Fig1: Total sugars (TS) and reducing sugars (RS) as influenced by moisture stress and irrigation modules




Fig2: Non-reducing sugars (NRS) and total soluble proteins(TSP) under moisture stress and irrigation modules.

Interactive effects between genotype x irrigation were significant too. Cultivars MLM-19 and NLM-3 possessed highest protein content of 8.8 mg g⁻¹ DW under moisture stress. NLM-3 registered highest protein content with one irrigation and NLM80 with two irrigations. RLC-1 registered the lowest amount of soluble proteins under all the three moisture regimes (Fig.2). Leaf protein content gradually decreases under moisture deficit conditions (Roy *et al.*, 2009). Further, increased protease enzyme activity under reduced moisture reduces protein content. Active protein breakdown pathway leads to reduced protein content under water stress as reported by Sankar *et al.* (2007) in groundnut. Our results are supported by the findings of Shahraki *et al.* (2008) and Tohidi *et al.* (2011) in *B.napus* where protein content decreased under water stress.

Proline content: Highest content of proline was 0.87 mg g⁻¹ DW under moisture stress (I₀) in NPJ-79 with a slight decline of 0.82 mg g⁻¹ DW in PLM-4. Under restricted moisture NLM-80 had highest proline content (0.69 mg g⁻¹ DW) while PLM-4 accumulated highest proline (0.49 mg g⁻¹ DW) under

normal moisture regime (Fig.3). Proline, one of the most common and compatible osmolytes in water stressed plants and its metabolism has been studied mainly in response to osmotic stress as reported by Verbruggen and Hermans (2008). Proline has the ability to oppose oxidative stress, an important strategy to overcome adverse effects of moisture stress (Vendruscolo *et al.*, 2007). Further, this molecule acts as a signaling molecule in modulation of mitochondrial functions or can trigger specific gene expression that can be essentially important for recovery of plant from stress as reported by Szabados and Savoure (2009). Proline accumulation therefore is considered as a drought tolerance mechanism which gets activated due to loss of feedback inhibition of proline synthesis which in turns declines the proline oxidation. Proline accumulation hence provides a good screening of drought resistant cultivars under water deficit conditions (Rahdari *et al.*, 2012). Results of the current investigation are corroborated by the findings in *B. napus* by Nosrati *et al.* (2014) and in soybean by Amira and Qados (2014).

Correlation: Correlation studies revealed negative association of seed yield with proline (-0.403) under




Fig3: Effect of moisture stress and irrigation modules on Proline content of the cultivars.

Table 1: Correlation coefficients of biochemical traits with yield under moisture stress (below diagonal) and restricted moisture (above diagonal)

Parameters	Total Sugars	Reducing Sugars	Non reducing sugars	Total soluble proteins	Proline	DSI ₁	DTI ₁	DSI ₂	DTI ₂	Seed yield
	1	2	3	4	5	6	7	8	9	10
Total sugars	1	-0.201	0.989**	-0.256	-0.321	-0.17	-0.171	-0.019	-0.099	0.045
Reducing sugars	-0.103	1	-0.341	-0.568	-0.171	0.422	0.425	0.417	0.115	0.39
Non reducing sugars	0.991**	-0.234	1	-0.162	-0.283	-0.227	-0.228	-0.081	-0.112	-0.015
Total soluble proteins	-0.117	-0.442	-0.055	1	0.527	-0.152	-0.178	0.087	0.002	-0.373
Proline	0.099	0.388	0.046	0.211	1	0.321	0.306	0.217	0.398	-0.253
DSI ₁	-0.219	0.253	-0.248	0.276	0.1	1	0.998**	0.652*	0.618*	0.371
DTI ₁	0.179	0.017	0.173	-0.172	-0.414	-0.124	1	.645*	0.618*	0.377
DSI ₂	-0.059	0.265	-0.093	0.38	-0.114	.652*	0.224	1	-0.046	0.377
DTI ₂	0.237	-0.022	0.235	-0.412	-0.203	-0.234	.845**	-0.174	1	-0.315
Seed yield	0.248	-0.099	0.256	-0.258	-0.403	-0.525	.909**	-0.077	.820**	1

DSI 1 and DTI 1: Seed yield at moisture stress (I₀) and restricted moisture (I₁)

DSI 2 and DTI 2: Seed yield at moisture stress (I₀) and normal moisture (I₂).

Table2: Correlation coefficients of biochemical traits with yield under normal moisture regime.

Parameters	Total sugars (TS)	Reducing sugars (RS)	Non-reducing sugars (NRS)	Total soluble proteins (TSP)	Proline	DSI ₁	DTI ₁	DSI ₂	DTI ₂	Seed yield
1	1	2	3	4	5	6	7	8	9	10
Total sugars	1									
Reducing sugars	0.26	1								
Non-reducing sugars		0.968**	0.143	1						
Total soluble proteins		-0.406	-0.398	-0.385	1					
Proline		0.262	0.157	0.241	1					
DSI ₁		0.046	-0.204	-0.248	0.162	1				
DTI ₁		0.029	-0.206	-0.275	0.148	.998**	1			
DSI ₂		0.304	-0.127	-0.396	0.207	.652*	.645*	1		
DTI ₂		-0.081	-0.036	0.248	0.276	.618*	.601*	-0.046	1	
Seed yield		0.145	0.091	0.143	-0.012	0.364	0.342	-0.179	0.626*	1

DSI 1 and DTI 1: Seed yield at moisture stress (I₀) and restricted moisture (I₁)DSI 2 and DTI 2: Seed yield at moisture stress (I₀) and normal moisture (I₂).

water stress which narrowed down to (-0.253) under restricted moisture and (-0.012) with normal irrigations (table 1 & 2). Thus, indicating higher amount of osmolytes as an adaptive strategy of plants to cope with water stress. Seed yield was positively and significantly correlated with DTI₁ (0.909**) and also DTI₂ (0.820**) under stress and only negatively related with DTI₂ (-0.315) with restricted moisture regime. Positive association existed between seed yield and DTI₂ (0.628*), DTI₁ and DSI₁ (0.988**), DSI₂ and DTI₁ (0.645*) and DTI₂ and DSI₁ (0.618*) with two irrigations.

Conclusion

Overall, water stress lead to increased accumulation of sugars and proline whereas total soluble protein content was reduced in the cultivars. The sugars and proline content in the cultivars was highest under moisture stress, followed by restricted moisture while least was under normal moisture (I₂). The soluble proteins were lowest under water stress and increased substantially with number of irrigation applied. K-9-108, NLM-3 and PLM-2 out yielded other cultivars under moisture stress due to high sugar levels and relatively higher proline content.

References

Ahmadi A, Emam Y and Pessarakli M. 2010. Biochemical changes in maize seedlings exposed to drought stress conditions at different nitrogen levels. *J Plant Nutr* **33**: 541-556.

Amira MS and Qados A. 2014. Effect of Ascorbic acid antioxidant on soybean (*Glycine max* L.) plants grown under water stress conditions. *Intl J Adv Res Biol Sci* **1**: 189-205.

Bates LS, Waldren RP and Teare ID. 1973. Rapid determination of free proline for water stress studies. *Plant Soil* **39**: 205-207.

Bayoumi TY, Eid MH and Metwali EM. 2008. Application of physiological and biochemical indices as a screening technique for drought tolerance in wheat genotypes. *African J Biotechnol* **7**: 2341-2352.

Dubois M, Gilles KA, Hamilton JK, Roberts PA and Smith F. 1956. Colorimetric methods for the determination of sugars and related substances. *Analyt Chem* **28**: 350-356.

Homayouni H and Khazarian V. 2014. Effects of deficit irrigation on soluble sugars, starch and

proline in three corn hybrids. *Indian J Sci Res* **7**: 910-917.

Jaleel CA, Gopi R and Panneerselvam R. 2008. Growth and photosynthetic pigments responses of two varieties of *Catharanthus roseus* to triadimefon treatment. *Comp Rend Biol* **31**: 272-277.

Lowry OH, Rosenbrough NJ, Farr AL and Randall RJ. 1951. Protein measurement with folin phenol reagent. *J Biol Chem* **193**: 265-275.

Mohammadkhani N and Heidari R. 2008. Effects of drought stress on soluble proteins in two maize varieties. *Turk J Bol* **32**: 23-30.

Nelson. 1944. A photometric adaption of the Somogyi method for the determination of glucose. *J Biol Chem* **153**: 375-380.

Nosrati S, Zanjan MJ and Asli DE. 2014. Fluctuations of proline concentration and soluble sugars content affected drought stress in canola (*Brassica napus* L.) seedlings. *J App Sci Agri* **9**: 497-502.

Praveen K, Singh RP and Singh C. 1996. Influence of irrigation scheduling on growth and seed quality of mustard. *Ann Agric Res* **17**: 184-85.

Rahdari P, Hosseini SM and Tavakoli S. 2012. The study effect of drought stress on germination, proline, sugar, lipid, protein and chlorophyll content in purslane (*Portulaca oleracea* L.) leaves. *J Med Plants Res* **6**: 1539-1547.

Roy, R, Mazumder, PB and Sharma, GD. 2009. Proline, catalase and root traits as indices of drought resistance in bold grained rice (*Oryza sativa*) genotypes. *African J Biotech* **8**: 6521-6528.

Sankar B, Jaleel CA, Manivannan P, Kishorekumar A, Somasundaram R and Panneerselvam R. 2007. Effect of paclobutrazol on moisture stress amelioration through antioxidants and free radical scavenging enzymes in *Arachis hypogaea* L. *Colloids Surf B Biointerfaces* **60**: 229-233.

Shahraki D, Nadian H, Bakhshandeh A, Fathi, Alamisaied K and Gharineh M. 2008. Optimization of irrigation and nitrogen regimes for rapeseed production under drought stress. *J Agron* **7**: 321-326.

- Sinay H and Karuwal RS. 2014. Proline and total soluble sugar content at the vegetative phase of six corn cultivars from Kisar island Maluku, grown under drought stress conditions. *Int J Adv Agri Res* **2**: 77-82.
- Szabados L and Savoure A. 2009. Proline: a multifunctional amino acid. *Trends Plant Sci* **15**: 89-97.
- Tahir M, Ali A, Nadeem MA, Tanveer A and Sabir QM. 2007. Performance of canola (*Brassica napus* L.) under different irrigation levels. *Pak J Bot* **39**: 739-746.
- Tohidi MHR, Farshad G and Zahedi H. 2011. Effect of UV radiation and elevated CO₂ on morphological traits, yield and yield components of canola (*Brassica napus* L.) grown under water deficit stress: *Not Bot Hort Agrobot Cluj* **39**: 213-219.
- Vendruscolo ACG, Schuster I, Pileggi M, Scapim CA, Molinari HBC, Marur CJ and Vieira LGC. 2007. Stress-induced synthesis of proline confers tolerance to water deficit in transgenic wheat. *J Plant Physiol* **164**: 1367-76.
- Verbruggen N and Hermans C. 2008. Proline accumulation in plants: a review. *Amino Acids* **35**: 753-759.
- Westgate ME and Boyer SS. 1985. Osmotic adjustment and the inhibition of leaf, root, stem and silk growth at low water potential in maize (*Zea mays* L.) *Planta* **164**: 540.
- Yordonov I, Velikova V and Tsonev T. 2003. Plant responses to drought and stress tolerance. *Bulg J Plant Physiol (Special issue)*: 187-286.
- Yunca H and Schmidhalter U. 2005. Drought and salinity. A comparison of the effects of drought and salinity. *J Plant Nutr Soil Sci* **168**: 541-49