



Performance of Gobhi season (*Brassica napus* L.) as influenced by different date of sowing and nitrogen levels under irrigated condition of central Punjab

Kamaldeep Kaur, Santosh Kumar* and Vivek Bhagat

Department of Agriculture, Mata Gujri College, Fatehgarh Sahib 140406, Punjab

*Corresponding author: santoshagro.nd@gmail.com

(Received: 23 February 2022; Revised: 20 June 2022; Accepted: 28 June 2022)

Abstract

A field experiment was carried out to assess the effects of different date of sowing and nitrogen levels on growth and yield of Gobhi sarson (*Brassica napus* L.) during *rabi* season of 2018-19. The experiment was laid out in split plot design with three replications with the three date of sowing as main plot treatment *viz.*, 10th October, 25th October and 15th November and four levels of N as sub-plot treatments *viz.*, 0, 40, 80 and 120 kg/ha. The results revealed that highest values of growth parameters, yield attributes and yield of crop were recorded under 25th October date of sowing which was remained at par with 10th October but it was recorded significantly superior over the 15th November. In case of N levels, the maximum values of growth parameters, yield attributes and yield of crop were recorded under 120 kg N/ha which was found at par with 80 kg N/ha but significantly superior over the other N levels. Therefore, 25th October sowing with 120 kg N/ha can be beneficial in enhancing the growth and yield attributes of the Gobhi sarson.

Keywords : Date of sowing, Gobhi sarson, N levels, yield attributes, yield

Introduction

India is one among the leading oil seed producing countries in the world. Oilseed brassica shares 24.4 % area and 26.8 % production of total oilseeds in the country (Choudhary *et al.*, 2019). It contributes more than 33 % of vegetable oil production (Langadi *et al.*, 2021). Gobhi sarson (*Brassica napus* L.) is the new emerging oilseed crops having limited area of cultivation. Sowing date is one of the most important agronomic factor and nonmonetary input which pave the way for better use of time and play an important role to fully exploit the genetic potentiality of a variety as it provides optimum growth conditions such as temperature, light, humidity and rainfall (Irradi, 2008; Singh *et al.*, 2019). Optimum time of sowing can provide congenial conditions to have maximum light interception, best utilization of moisture and nutrients from early growth stage to seed filling stage (Keerthi *et al.*, 2017). The significant reduction in yield of mustard due to delay in sowing of crop from September to December has been reported by many workers (Khushu and Singh, 2005; Alam *et al.*, 2014). Further, Kumar *et al.* (2018) also reported that the values of yield and yield attributes were highest on 16th October sowing while 21st November sowing dates showed lowest values of yield and yield attributing traits. Similarly, the maximum seed yield was recorded when crop sown on 25th October followed by 15th October sowing (Jat *et al.*, 2019).

Furthermore, plant nutrition is a key input to increase the productivity of mustard seed crop. Nitrogen (N) is the most important nutrient, for the crop to activate the metabolic activity and transformation of energy, chlorophyll and protein synthesis and being a constituent of protoplasm and protein (Bhattacharya, 2014). When nitrogen supplies are optimum and conditions are favourable for growth, proteins are formed from the manufactured carbohydrates. Nitrogen also affects uptake of other essential nutrients and it helps in the better partitioning of photosynthates to reproductive parts which increase the seed: stover ratio and enhances the yield of rapeseed mustard group of crops (Singh and Meena, 2004). Significantly higher values of primary as well as secondary branches, siliquae/plant and seed yield of Indian mustard was reported with the application of N up to 120 kg/ha (Yadav *et al.*, 2007). The plant height, leaf area index and dry matter production were significant increased with increasing levels of N rates from 40 to 120 kg/ha (Rasool *et al.*, 2013). Hence, identification of a suitable combination of date of sowing and nitrogen level could be helpful in maximizing the yield of Gobhi sarson in the Punjab.

Materials and Methods

A field experiment was conducted at experimental farm, Mata Gujri College, Fatehgarh Sahib during *Rabi* season of year 2018-19. The experiment was laid out in split plot

Crop yield and harvest index

The data pertaining to seed yield, stover yield and biological yield of Gobhi sarson were significantly influenced by application of different treatments. However, harvest index was not significantly influenced by different treatments (Table 3). From the data it is evident that the maximum seed yield, stover yield and biological yield were observed in 25th October sowing which was at par with 10th October and which was significantly superior over the 15th November. Further, it was higher 4.18 %, 4.23 % and 4.23 % to 10th October sowing and 11.33 %, 12.09 % and 11.92 % to 15th November sowing in seed yield, stover yield and biological yield, respectively. Because in optimum sowing time the crop may be ascribed mainly to prolonged duration of reproductive period indicated more time available for utilization of assimilates for seed setting which led to increase seed yield. The biological yield also increases because there is positive and significant correlation also existed between stover and biological yield. Similar findings have also been reported by (Keerathi *et al.* 2017). Among nitrogen application, the maximum seed yield, stover yield and biological yield were observed in 120 kg N/ha which was statistically at par with 80 kg N/ha and which was significantly superior over other treatments. It is higher 3.52 %, 3.96 % and 3.86 % to 80 kg N/ha and 19.9 %, 9.12 and 11.62 % to 0 kg N/ha in seed yield, stover yield and biological yield, respectively. It is due to that the increase in stover yield with nitrogen application may attribute to favorable effect of nitrogen growth parameters and yield attributes of crop with application of higher doses of nitrogen over lower doses accounted for increased seed yield. Interaction effect of date of sowing and nitrogen application were do not affect significantly on seed yield, stover yield and biological yield. Similar results have been reported by Kaur *et al.* (2018). In case of harvest index, the maximum harvest index was observed in 15th November sowing which was closely followed by 25th October sowing of crop. It is due to the good supply of nutrition which ultimately increases in harvest index. Similar results have been given by Kaur *et al.* (2018). Under nitrogen effect, maximum harvest index was observed in 80 kg N/ha which was closely followed by 120 kg N/ha while minimum was observed in 0 kg N/ha during experimentation.

Conclusion

It can be concluded that growth contributing parameters and yield attributes of Gobhi sarson were positively influenced due to date of sowing and nitrogen levels. Hence, the crop sown on 20th October with nitrogen 120 kg/ha resulted the better growth parameters, yield

attributing traits and yield of Gobhi sarson under irrigated conditions of central Punjab.

References

- Alam MM, Begum F and Roy P. 2014. Yield and yield attributes of rapeseed-mustard (*Brassica* spp.) genotypes grown under late sown condition. *Bangladesh J Agric Res* **39**: 311-336.
- Bhagat R, Singh M, Sharma BC and Shankar U. 2022. Effect of sowing environments on yield attributes and yield of Gobhi Sarson (*B. napus*) cultivars under sub-tropics of Jammu region. *J Oilseed Brassica* **13**(1): 7-10.
- Bhattacharya G. 2014. Soil fertility and nutrient management. Oxford Book Company, New Delhi, pp 109-110.
- Choudhary RL, Langadi AK, Jat RS, Anupama, Singh HV, Meena MD, Dotaniya ML, Meena MK, Premi OP and Rai PK. 2019. Mitigating the moisture stress in Indian mustard (*B. juncea*) through polymer. *J Oilseed Brassica* **12**: 21–27.
- Iraddi VS and Mansur CP. 2008. Response of mustard (*B. juncea*) varieties to date of sowing and row spacing in Northern transition zone of Karnataka. Thesis Submitted to the University of Agricultural Sciences, Dharwad in partial fulfilment of the requirements for the Degree of Master of Science (Agriculture) in Agronomy, pp 51-81.
- Jat AL, Desai AG and Rathore BS. 2019. Effect of different sowing schedule and crop geometry on productivity and profitability of Indian mustard (*Brassica juncea* L.). *J Oilseed Res* **36**: 17-19.
- Kaur L, Sardana V and Sharma P. 2018. Effect of sowing dates and nitrogen application on growth and productivity of canola oilseed rape (*Brassica napus*). *J Oilseed Brassica* **9**: 114-121.
- Keerthi P, Pannu RK, Dhaka AK and Chaudhary K. 2017. Effect of sowing time and nitrogen on growth, yield and nutrient uptake by Indian mustard (*B. juncea*) under Western Haryana. *Chem Sci Rev Letter* **6**: 2526-2532.
- Khushu M and Singh M. 2005. Thermal response of mustard under rainfed condition of Jammu. *Environ Ecol* **23**: 683-686.
- Kumar A, Lal M, Mohan N, Kumar M and Kumar N. 2018. Effect of different sowing dates on yield and yield attributes of Indian Mustard (*B. juncea*) Genotypes. *Int J Pure Applied Biosci* **6**: 848-853.
- Kumar S, Sairam RK and Prabhu KV. 2013. Physiological traits for high temperature stress tolerance in *Brassica juncea*. *Indian J Plant Physiol* **18**: 89-93.

- Langadi AK, Choudhary RL, Jat RS, Singh HV, Dotaniya ML, Meena MK, Premi OP and Rai PK. 2021. Effect of superabsorbent polymer on drought Mitigation, and enhancing productivity and profitability of Indian mustard (*B. juncea*). *J Oilseeds Res* **38**: 179–186.
- Panse VG and Sukhatme PV. 1978. Statistical Methods for Agricultural Workers. 2nd Ed. Indian Council of Agricultural Research, New Delhi, India.
- Rasool F, Hasan B, Jahangiri IA, Ali T and Mubarak T. 2013. Nutritional yield and economic responses of sunflower (*H. annuus*) to integrated levels of nitrogen, sulphur and farm yard manure. *J Agric Sci* **8**: 17-27.
- Singh A and Meena NL. 2004. Effect of nitrogen and sulphur on growth, yield attributes and seed yield of mustard (*B. juncea*) in eastern plains of Rajasthan. *Indian J Agron* **49**: 186-188.
- Singh AK, Singh RR, Singh AK and Singh PK. 2014. Influence of dates of sowing and irrigation scheduling on growth and yield of mustard (*B. juncea*). *Int J Farm Sci* **4**: 80-85.
- Singh HV, Choudhary RL, Meena MD, Dotaniya ML, Meena MK, Jat RS, Premi OP and Rai PK. 2019. Productivity enhancement of Indian mustard (*B. juncea*) through innovative crop establishment technique. *Indian J Agric Sci* **89**: 1669–73.
- Yadav RB, Singh RK and Singh H. 2007. Response of Indian mustard (*B. juncea*) to nitrogen and sulphur in mid-western plain zone of Uttar Pradesh. *Indian J Crop Sci* **2**: 243-44.