



Trait modelling for stress tolerance in Indian mustard : Evidenced from seedling stage

G Kumawat¹, ML Jakhar¹, V Singh, DK Gothwal¹ and J Singh*

ICAR-Central Soil Salinity Research Institute, Karnal 132001, Haryana, India

¹Sri Karan Narendra Agriculture University, Jobner 303329, Jaipur, Rajasthan, India

*Corresponding author: jogendra.singh@icar.gov.in

(Received: 17 December 2022; Revised: 31 December 2022; Accepted: 05 January 2023)

Abstract

The current scanty knowledge about the physiological mechanisms underlying plants' ability to tolerate salt stress that hinders potential production of numerous crops, including mustard. To explore the traits and mechanism for salt tolerance in mustard, we used 250 stabilized $F_{7,8}$ recombinant inbred lines (RILs) mapping population developed by crossing indigenous contrasting genotypes CS 614-1-1-100-13 (salt sensitive) \times CS 56 (salt tolerant) and evaluated them under control and irrigated water salinity of EC_{iw} 12 dS/m to characterize for growth, photosynthetic and ionic traits. Step wise regression revealed, instantaneous water use efficiency, transpiration rate and fresh weight of root together accounted for more than 93% of the overall variation in photosynthesis rate under salt stress condition, indicating their critical contribution to reducing salt stress. The salt tolerance index (STI) categorized 23 RILs as salt tolerant, 99 RILs as moderately tolerant and remaining 128 RILs were sorted as salt sensitive RILs. These identified salt tolerant RILs can be exploited for QTLs and gene discovery and serve as potential donors/future ready lines to combat abiotic stress and development of salt tolerant varieties of mustard.

Keywords: Mustard, RILs, photosynthetic traits, STI

Introduction

Salinity and sodicity stress damage an area of 932.2 mha worldwide, of which 6.73 mha are affected by these stresses in India (Metternicht and Zinck, 2003; Singh *et al.*, 2014) however, this will be increased up to 16.25 mha by 2050 (Kumar *et al.*, 2022b). Groundwater utilized for irrigation contains 32-84 percent salty or brackish water and the soil salinity build due to this irrigation results in an annual loss of 10 million ha of land, consequently adverse effect on the food basket. High salt stress leads to cellular osmotic stress, ion-specific toxicity, reduced plant growth and photosynthetic traits which ultimately lead to a minimal yield of the crop (Singh and Sharma, 2016). By improving crop tolerance to salt or by draining salt from the soil, this low yield in saline places can be combated (Kumar, 2014; Singh and Sharma, 2016). This tolerance is achieved in crop plants *via* different mechanisms such as maintenance of cell turgidity through the accumulation of osmolytes, ion exclusion from the root, ion compartmentation in vacuole, tissue tolerance and ion-independent tolerance.

Indian mustard, *Brassica juncea* (L.) Czern & Coss (AABB, $2n=36$, Genome size: 1068 Mb) is a significant oilseed crop with wide adaptability (Kang *et al.*, 2021). Globally,

India ranked second in rapeseed-mustard cultivation after only China and third in production behind Canada and China (Kumari *et al.*, 2019). It ranks as India's third-largest edible oilseed crop after peanut and soybean, which accounts for 24.36% of the country's oilseed market among nine edible oilseed crop (Kumar *et al.*, 2022a). The amphidiploid species (*B. carinata*, *B. juncea* and *B. napus*) appear to be superior to the diploid species (*B. rapa*, *B. nigra* and *B. oleracea*) in terms of saline tolerance, as per several studies (Ashraf *et al.*, 2001; Ashraf and Mehmood, 1990).

Salt stress causes Na^+ and K^+ ion imbalance, by disrupting the Na^+/K^+ ratio in leaves, salinity during the seedling stage has a deleterious impact on photosynthesis. The transfer of carbohydrates from source to sink is slowed down by this aberrant Na^+/K^+ ratio, which also affects mustard growth. Due to restrictions on growth, seed yield is reduced by up to 60% in mustard (Singh *et al.*, 2019). The lowered rate of carbon absorption, assimilation and partitioning to seedlings during the initial stage of salt stress makes it more detrimental (Singh *et al.*, 2019). Ultimately, plant growth, photosynthesis rate and yield is reduced due to toxic salt stress levels (Pant *et al.*, 2022).

It's well-recognized that *Brassica* an acceptable

reclamation crop but introduction of salt mitigation or salt tolerance mechanisms into the Indian mustard crop proceeds slowly due to a lack of genetic variability, research or inadequate information in these areas. One of the main initiatives in *Brassica* development is cultivar improvement for saline tolerance. Therefore, the creation of salinity-tolerant Indian mustard cultivars with higher yields in the salt-affected semi-arid tropics will be more effective and efficient by integrating breeding, physiological, biochemical, novel omics as well as bioinformatics approaches studies (Pant *et al.*, 2022). Our study included 250 RILs mapping population which were produced by crossing indigenous contrasting genotypes CS 614-1-1-100-13 (salt sensitive) \times CS 56 (salt tolerant) of *B. juncea*. These RIL mapping population serve as mustard genetic resource to identify salt tolerance genotypes to combat salt stress for salt prone areas.

Materials and Methods

Study site

The experimental materials consisted of 250 stabilized $F_{7:8}$ recombinant inbred lines (RILs) mapping population of Indian mustard developed by crossing indigenous sources of mustard; CS 614-1-1-100-13 (salt sensitive) an advanced breeding line developed at ICAR-CSSRI, Karnal with gamma-ray irradiation treatment and stabilized for M_6 generation (Sharma *et al.*, 2008) and CS 56, a national released high yielding salt tolerant variety. These 250 RILs along with parents grown during consecutive *Rabi* seasons 2020-21 and 2021-22 under control and irrigation water salinity EC_{iw} 12 dS/m in the pots with three replications at ICAR-CSSRI, Karnal ($29^{\circ}43'N$, $76^{\circ}58'E$; 245 m above the average sea level) (Singh *et al.*, 2020) (Fig. 1).



Fig. 1: Location of experimental site

Experimental details

The RIL mapping population was cultivated in pots of 20 kg capacity in sand culture. For the salinity environment, irrigated with saline water of EC_{iw} 12 dS/m throughout the experiment. The chloride and sulphate salts of Na^+ , Ca^{2+} and Mg^{2+} to keep the SAR (Sodium absorption ratio)

within the permissible limits used for the preparation of EC_{iw} 12 dS/m saline irrigation water. Prior to planting, seeds were surface sterilised for 5 minutes in a solution of 10% sodium hypochlorite before being rinsed with distilled water. Twenty seeds of each RIL were planted in a plastic pot filled with properly washed river sand at a depth of one centimetre. Each pot's bottom was dug out to allow any extra water to drain. The pots were set up in a factorial experiment using a completely random block design. The pots were watered with Hoagland's solution, a nutrient solution and kept at maximum field capacity until the seedling stage. Throughout the experiment, salinity levels were kept constant by draining the salt out of the pots every day (Singh *et al.*, 2019).

Data collection

Growth attributing traits at the seedling stage

Initially, randomly ten seedlings (15 days old) from each RIL under control and salinity EC_{iw} 12 dS/m conditions were uprooted and washed with distilled water to record the fresh weight (mg) of leaves, stems and roots. The oven-dried plant samples at $55-65^{\circ}C$ for 5-6 days were used for reading of dry weight (g) of the leaf, stem and root.

Photosynthetic traits

Randomly selected three plants from each genotype under control and EC_{iw} 12 dS/m salinity regime were used for photosynthetic data at the seedling stage *i.e.*, Photosynthesis rate (P_n ; $\mu\text{mol m}^{-2} \text{s}^{-1}$), transpiration rate (E ; $\text{mol m}^{-2} \text{s}^{-1}$), stomatal conductance (g_s ; $\text{mol m}^{-2} \text{s}^{-1}$), intracellular CO_2 assimilation (C_i/C_a), Instantaneous water use efficiency ($iWUE$; $\mu\text{mol } CO_2/\text{m mol } H_2O$) and intrinsic water use efficiency ($inWUE$). P_n and other gas exchange parameters were measured on the fully expanded leaf of three representative plants per RILs using a portable photosynthetic system: infrared gas analyzer LI-6800XT [Li-COR, USA] (Singh *et al.*, 2019). All the above traits were measured between 10:00 and 12:00 AM in sunlight under these weather conditions; PAR $\sim 700 \mu\text{mol m}^{-2} \text{s}^{-1}$, temperature $\sim 25 \pm 1^{\circ}C$ relative humidity $\sim 70\%$ and air CO_2 $355 \mu\text{mol mol}^{-1}$. The K^+ and Na^+ concentration of plant sample measured with Inductively coupled plasma-optical emission spectrometry ICPE-9800 (Shimadzu, Japan) after checking the standards (Piper, 2019).

Statistical analysis

The statistical analyses regression and STI was carried out for all the studied seedling stage traits using the *STAR 2.0.1* (IRRI, 2014) and MS Excel.

$$\text{Stress tolerance index (Fernandez, 1992)} = \frac{Y_s Y_p}{Y_p^2}$$

Where, Y_s and Y_p are the mean yield of genotypes under stress and non-stress conditions, respectively. The RILs with high STI values will be salt stress tolerance Result and discussion

Twenty-one traits were further sub-categorized into six growth attributing traits [fresh weight (root, stem and leaves), dry weight (root, stem and leaves)], six photosynthetic traits and nine ionic traits (Na^+ , K^+ and Na^+/K^+ of root, stem and leaves) were used for trait modeling and characterized salt tolerant RILs at the seedling stage.

Results and Discussion

Mustard traits prioritization under salinity

All feasible and stepwise regression analyses were carried out to ascertain the impact of component variables on Pn (dependent variable) (Shannon *et al.*, 2000; Sharma and Sinha, 2012). All conceivable regression analyses revealed that gS, E, *i*WUE, *in*WUE, RFW and LFW significantly influenced the Pn of mustard leaves under salt stress, while the remaining traits did not (Table 1). Therefore, during the stepwise regression method, these remaining non-significant traits were eliminated. According to the results, *i*WUE, E and RFW together accounted for more than 93% of the overall variation in Pn under salt stress conditions. Additionally, Pn variation was substantially

influenced by *i*WUE, E, RFW, *in*WUE, gS and LFW with cumulative $R^2 = 94.36$, which could be best fitted since it reflected the least Mallows' Cp criteria. The following equation was created to estimate the projected Pn under saline conditions based on regression coefficients of the relevant traits (Table 2):

$$\text{Predicted photosynthetic rate} = -17.05 + (3.8 \times E) + (gS \times 13.13) + (2.44 \times iWUE) + (0.08 \times inWUE) + (-0.02 \times RFW) + (-0.01 \times SFW) + (0.23 \times LFW) + (0.01 \times LK15).$$

Hence, these traits play major in contribution to enhance photosynthetic rate under saline condition. By targeting these traits, further research will be extending to vegetative and harvesting stage to combat salt stress.

Characterization of RILs based on salt tolerance index

To characterized tolerant and sensitive RILs over the environment, the salt tolerance index (STI) was calculated. Based on STI, RILs were characterized into three groups *i.e.*, highly tolerant (STI ≥ 1), tolerant (STI = 1-0.75), moderately tolerant (STI = 0.75-0.50) and sensitive RILs (STI < 0.50). Total 24 RILs were selected as highly salt tolerant (RIL24, RIL32, RIL74, RIL87 and RIL73), 99 RILs were sorted as tolerant (RIL14, RIL31, RIL13, RIL95 and RIL142), 109 RILs as moderately tolerant while remaining 18 RILs were characterized as sensitive which score STI < 0.50 (RIL170, RIL247, RIL225, RIL228 and RIL223) over the environment (Table 3 and Fig. 2). The salt-tolerant

Table 1: Salinity stress tolerance's regression coefficient, standard error and significance of the prioritized attributes

Dependable variable	Variable	Estimate	Standard Error (SE)	t value	Pr(> t)
Pn	Intercept	-17.05	0.71	-23.97	0.000
	E	3.80	0.19	19.60	0.000
	gS	13.13	1.95	6.72	0.000
	<i>i</i> WUE	2.44	0.13	18.93	0.000
	<i>in</i> WUE	0.08	0.01	6.66	0.000
	RFW	-0.02	0.01	-3.85	0.000
	SFW	-0.01	0.00	-1.53	0.127
	LFW	0.23	0.14	1.72	0.087
	LK15	0.01	0.00	1.78	0.077

Table 2: Traits modelling for salinity tolerance through multiple linear regressions approach

Variables	C(p)	R-square	Adj R-sq
<i>i</i> WUE	2845.40	29.19	28.90
E + <i>i</i> WUE	56.81	93.07	93.01
E + <i>i</i> WUE + RFW15DAS	47.63	93.32	93.24
GSW + E + <i>i</i> WUE + <i>in</i> WUE	19.57	94.01	93.92
GSW + E + <i>i</i> WUE + <i>in</i> WUE + RFW15DAS	8.49	94.31	94.20
GSW + E + <i>i</i> WUE + <i>in</i> WUE + RFW15DAS + LFW15DAS	8.40	94.36	94.22

Table 3: Grouping of mustard RILs based on salt tolerance index (STI) of photosynthetic rate

Highly tolerant RILs (STI > 1)			Tolerant RILs (STI = 1-0.75)			Moderately tolerant RILs (STI = 0.75-0.50)			Sensitive RILs (STI = <0.50)		
RILs	STI	Rank	RILs	STI	Rank	RILs	STI	Rank	RILs	STI	Rank
RIL24	1.26	1	RIL14	0.99	25	RIL128	0.74	124	RIL169	0.64	179
RIL32	1.22	2	RIL31	0.99	26	RIL141	0.85	76	RIL187	0.64	180
RIL74	1.18	3	RIL13	0.99	27	RIL111	0.84	77	RIL65	0.64	181
RIL87	1.17	4	RIL95	0.97	28	RIL97	0.84	78	RIL154	0.64	182
RIL73	1.15	5	RIL142	0.97	29	RIL34	0.84	79	RIL61	0.64	183
RIL77	1.15	6	RIL78	0.97	30	RIL96	0.84	80	RIL148	0.63	184
RIL25	1.13	7	RIL56	0.97	31	RIL174	0.84	81	RIL113	0.63	185
RIL82	1.12	8	RIL161	0.96	32	RIL88	0.84	82	RIL221	0.63	186
RIL30	1.12	9	RIL102	0.96	33	RIL147	0.84	83	RIL133	0.63	187
RIL79	1.1	10	RIL51	0.96	34	RIL29	0.84	84	RIL237	0.63	188
RIL22	1.1	11	RIL49	0.95	35	RIL17	0.83	85	RIL136	0.62	189
RIL37	1.07	12	RIL41	0.95	36	RIL149	0.83	86	RIL240	0.62	190
RIL83	1.06	13	RIL108	0.95	37	RIL19	0.83	87	RIL235	0.62	191
RIL117	1.04	14	RIL26	0.94	38	RIL110	0.83	88	RIL239	0.62	192
RIL3	1.03	15	RIL57	0.94	39	RIL6	0.83	89	RIL168	0.62	193
RIL46	1.03	16	RIL242	0.94	40	RIL5	0.83	90	RIL158	0.62	194
RIL99	1.02	17	RIL129	0.94	41	RIL47	0.83	91	RIL81	0.61	195
RIL36	1.02	18	RIL93	0.93	42	RIL241	0.82	92	RIL216	0.61	196
RIL94	1.01	19	RIL85	0.93	43	RIL220	0.82	93	RIL11	0.61	197
RIL109	1.01	20	RIL91	0.93	44	RIL90	0.82	94	RIL206	0.61	198
RIL72	1.01	21	RIL100	0.93	45	RIL209	0.81	95	RIL153	0.61	199
RIL107	1.01	22	RIL131	0.92	46	RIL101	0.81	96	RIL28	0.6	200
RIL76	1.00	23	RIL27	0.92	47	RIL39	0.81	97	RIL214	0.6	201
RIL33	1.00	24	RIL68	0.92	48	RIL16	0.81	98	RIL62	0.59	202
			RIL197	0.91	49	RIL115	0.81	99	RIL90	0.59	203
			RIL145	0.91	50	RIL42	0.8	100	RIL137	0.59	204
			RIL69	0.91	51	RIL159	0.8	101	RIL156	0.59	205
			RIL71	0.91	52	RIL84	0.8	102	RIL162	0.59	206
			RIL132	0.91	53	RIL196	0.8	103	RIL163	0.59	207
			RIL160	0.91	54	RIL125	0.8	104	RIL165	0.58	208
			RIL23	0.91	55	RIL218	0.8	105	RIL238	0.56	209
			RIL53	0.9	56	RIL127	0.8	106	RIL213	0.56	210
			RIL12	0.9	57	RIL59	0.8	107	RIL224	0.56	211
			RIL140	0.9	58	RIL130	0.8	108	RIL236	0.55	212

RIL86	0.9	59	RIL191	0.79	109	RIL188	0.69	158	RIL229	0.55	213
RIL55	0.9	60	RIL146	0.79	110	RIL171	0.69	159	RIL244	0.54	214
RIL48	0.89	61	RIL150	0.79	111	RIL234	0.69	160	RIL202	0.54	215
RIL63	0.89	62	RIL50	0.79	112	RIL44	0.69	161	RIL164	0.54	216
RIL126	0.89	63	RIL2	0.78	113	RIL192	0.68	162	RIL176	0.53	217
RIL104	0.89	64	RIL98	0.78	114	RIL20	0.68	163	RIL138	0.53	218
RIL139	0.89	65	RIL60	0.77	115	RIL144	0.68	164	RIL195	0.52	219
RIL103	0.87	66	RIL7	0.77	116	RIL9	0.67	165	RIL210	0.52	220
RIL122	0.87	67	RIL18	0.77	117	RIL215	0.67	166	RIL181	0.52	221
RIL106	0.87	68	RIL45	0.76	118	RIL198	0.67	167	RIL177	0.52	222
RIL116	0.86	69	RIL40	0.76	119	RIL66	0.67	168	RIL227	0.52	223
RIL208	0.86	70	RIL89	0.76	120	RIL64	0.66	169	RIL226	0.52	224
RIL15	0.86	71	RIL119	0.76	121	RIL134	0.66	170	RIL245	0.52	225
RIL75	0.85	72	RIL38	0.75	122	RIL193	0.66	171	RIL203	0.52	226
RIL58	0.85	73	RIL123	0.75	123	RIL219	0.65	172	RIL200	0.51	227
RIL80	0.85	74				RIL67	0.65	173	RIL173	0.51	228
						RIL232	0.65	174	RIL250	0.5	229
						RIL248	0.65	175	RIL243	0.5	230
						RIL212	0.65	176	RIL178	0.5	231
						RIL92	0.65	177	RIL199	0.5	232
						RIL207	0.65	178			




Fig. 2: Salinity tolerant and sensitive RILs of mustard under salinity environment EC_{iw} 12dS/m at seedling stage

RILs under salt stress circumstances due to the preservation of high photosynthetic activity, high K^+ concentration, low Na^+ and Na^+/K^+ ratio (Keisham *et al.*, 2018). These research' conclusions agreed with our experimental results.

A multiple regression model's fit can be evaluated using Mallows' Cp Criterion; smaller Cp values are preferable because they signify lower levels of unexplained error.

Conclusion

The Indian mustard RILs evaluated in our study have a significant variation in measured growth, photosynthetic and ionic traits. The stepwise regression approach $iWUE$, E , RFW, $inWUE$, gS and LFW as defining traits for Pn, indicating their critical contribution to reducing salt stress. Based on our study on RIL24, RIL32, RIL74 and RIL 87 were identified as potential resource or donor for salt-tolerance and may be used for cultivation under salinity stress. Further these lines may be employed in hybridization programs to create future ready new high-yielding, salt-tolerant breeding lines to combat salt stress. Additionally, these genotypes might be used to comprehend the genetic and molecular mechanism of Indian mustard salt tolerance.

References

Ashraf M and Mehmood S. 1990. Response of four *Brassica* species to drought stress. *Environ Exp Bot* **30**: 93–100. DOI: 10.1016/0098-8472(90)90013-T.

Ashraf M, Nazir N and McNeilly T. 2001. Comparative salt tolerance of amphidiploid and diploid *Brassica* species. *Plant Sci* **160**: 683–689.

Fernandez GCJ. 1992. Effective selection criteria for assessing plant stress tolerance. *Proceeding of the International Symposium on Adaptation of Vegetables and other Food Crops in Temperature and Water Stress, Shanhuia, Taiwan*, 1992: 257–270.

IRRI. 2014. Statistical Tool for Agricultural Research (STAR) v. 2.0. 1. IRRI Los Banos, Philippines.

Kang L, Qian L, Zheng M, Chen L, Chen H, Yang L, You L, Yang B, Yan M, Gu Y, Wang T, Schiessl SV, An H, Blischak P, Liu X, Lu H, Zhang D, Rao Y, Jia D, Zhou D, Xiao H, Wang Y, Xiong X, Mason AS, Chris Pires J, Snowdon RJ, Hua W and Liu Z. 2021. Genomic insights into the origin, domestication and diversification of *B. juncea*. *Nat Genet* **53**: 1392–1402. DOI: 10.1038/s41588-021-00922-y.

Keisham M, Mukherjee S and Bhatla SC. 2018. Mechanisms of sodium transport in plants—progresses and challenges. *Int J Mol Sci* **19**: 647.

Kumar M. 2014. Crop plants and abiotic stresses. *J Biomol Res Ther* **3**. DOI: 10.4172/2167-7956.1000e125.

Kumar PP, Rao SVR, Koshta VKCAK, Khan MA, Lakhera ML and Teja IK. 2022a. Performance of rapeseed-mustard in India—a temporal analysis. *J Oilseed Brassica* **13**: 45–52.

Kumar R, Singh A, Bhardwaj AK, Kumar A, Yadav RK and Sharma PC. 2022b. Reclamation of salt-affected soils in India: Progress, emerging challenges, and future strategies. *L Degrad Dev* **33**: 2169–2180. DOI: 10.1002/ldr.4320.

Kumari V, Jambhulka S, Sharma HKCBK, Sood P, Guleria SK, Bala A and Sanju S. 2019. Phenotypic stability for seed yield and related traits in Trombay mustard genotypes under North western Himalayas. *J Oilseed Brassica* **10**: 33–37.

Metternicht GI and Zinck JA. 2003. Remote sensing of soil salinity: Potentials and constraints. *Remote Sens Environ* **85**: 1–20. DOI: 10.1016/S0034-4257(02)00188-8.

Pant P, Hamsa S and Kaur J. 2022. Advances in Breeding Strategies for Improving Stress Tolerance in *Brassic*s. The *Brassica juncea* Genome. Springer, 439–469.

- Piper CS. 2019. Soil and plant analysis. *Scientific Publishers*.
- Shannon MC, Grieve CM, Lesch SM and Draper JH. 2000. Analysis of salt tolerance in nine leafy vegetables irrigated with saline drainage water. *J Am Soc Hortic Sci* **125**: 658–664.
- Sharma PC and Sinha TS. 2012. Salt tolerance of Indian mustard: physiological factors. *Int J Life Sci* **1**: 97–111.
- Sharma PC, Prashat R, Gingham GD and Pareek A. 2008. Improving salt tolerance and seed yield in Indian mustard (*B. juncea* L.) through radiation induced mutagenesis. *Int Symp Ind Mut Plant*: No. IAEA-CN—167.
- Singh A, Singh RK, Singh J, Goswami A, Upadhyaya A and Sharma PC. 2020. 'DUS' characterization of an endangered salt tolerant radish landrace (Newar). *Indian J Tradit Knowl* **19**: 24–32.
- Singh J and Sharma PC. 2016. Comparative effects of soil and water salinity on oil quality parameters of *B. juncea*. *J Oilseed Brassica* **7**: 29–37.
- Singh J, Sharma PC, Sharma SK and Rai M. 2014. Assessing the effect of salinity on the oil quality parameters of Indian mustard (*B. juncea*) using Fourier Transform Near-Infrared Reflectance (FT-NIR) spectroscopy. *Grasas y Aceites* **65**: 1–8.
- Singh J, Singh V, Vineeth T V., Kumar P, Kumar N and Sharma PC. 2019. Differential response of Indian mustard (*B. juncea*) under salinity: photosynthetic traits and gene expression. *Physiol Mol Biol Plants* **25**: 71–83.