



## Effect of *Sclerotinia sclerotiorum* culture filtrate on seed germination and seedling vigour of Indian mustard (*Brassica juncea* cv. Rohini)

Pankaj Sharma\*, PD Meena and Dhiraj Singh

Directorate of Rapeseed-Mustard Research (ICAR), Bharatpur 321 303, India

\*Corresponding author: pksvirus@gmail.com

(Received: 20 February 2014; Revised: 17 April 2014; Accepted: 27 May 2014)

### Abstract

Culture filtrate of 25 *Sclerotinia sclerotiorum* isolates isolated from infected *Brassica* plants grown in nine states in India were tested for their effect on germination and seedling vigour of Indian mustard (*Brassica juncea*) cultivar Rohini; fresh and dry mycelial weight of all isolates grown in Richard's liquid medium were also recorded. All geographical isolates showed considerable variability in fresh and dry mycelial weight. Compared to the control, filtrates of all isolates reduced seed germination and length of radicle and plumule. Isolates SR 15 and SR 03 significantly reduced the germination (53.5 and 53.3%) and gave the maximum pre-emergence mortality (46.9 and 46.7%). However, isolate SR 19 gave the maximum seed germination (83.4%) and minimum pre-emergence mortality (16.7%).

**Key words:** Culture filtrate, *Sclerotinia sclerotiorum*, seed germination, seedling vigour

### Introduction

*Sclerotinia sclerotiorum* (Lib.) de Bary is one of the most devastating and cosmopolitan fungal pathogen belonging to ascomycetes and capable of infecting more than 400 host plants in 75 families (Boland and Hall, 1994). In India, *Sclerotinia* rot is a threat to cultivation of oilseed Brassica in Rajasthan, Haryana and Punjab states of North India. The disease is important particularly in areas where mono-cropping is practiced. Disease incidence as high as 80 % has been recorded in some parts of Punjab, MP, UP and Haryana (Aggrawal *et al.*, 1997). A necrotrophic pathogen with cosmopolitan distribution and a wide host range, *S. sclerotiorum* shows a high level of intraspecific phenotypic variability (Purdy, 1979). Variability among *S. sclerotiorum* populations based generally on morphological differences in sclerotia, mycelial growth, and ascospores has also been reported (Attalah *et al.*, 2004; Harlton *et al.*, 1995; Okabe *et al.*, 1998). Fungal metabolites are substances discharged by fungi in their metabolic processes and are products of some amino acids, cyclic peptides, aromatic, phenols, terpenoids and plant growth regulators (Griffin, 1981; Madhosing, 1995). Culture

filtrate from different pathogenic fungi has been reported to reduce seed germination and seedling vigour. Reduction in % seed germination of soybean seeds was observed in seeds soaked in filtrates of *Phomopsis phaseoli* (Hilty and Lee, 1988). Filtrate from mycelial cultures of *Verticillium albo-atrum* was found to inhibit cell growth and reduced the viability of alfalfa (*Medicago sativa*) seeds (Frame *et al.*, 1991). Culture filtrates of *Fusarium moniliforme*, *F. semitectum* and *F. oxysporum* are also known to reduce seed germination and inhibit root and shoot growth of sorghum. The present study was therefore, undertaken to determine the effect of culture filtrates of different geographical isolates of *Sclerotinia sclerotiorum* from Brassica on seed germination and seedling vigour of Indian mustard.

### Materials and Methods

#### *Sclerotinia sclerotiorum* isolates

Twenty five *Sclerotinia sclerotiorum* isolates from infected Brassica plants were collected from Bihar, Himachal Pradesh, Jammu and Kashmir, Madhya Pradesh, Punjab, Rajasthan, UttaraKhand, Uttar Pradesh and West Bengal. *S. sclerotiorum* was isolated from infected plant tissues and designated

as SR-1 to SR- 25 (Table 1). For isolation, a single sclerotium or infected host tissue was surface-sterilized in 5% sodium hypochlorite solution for 5 min, in 70% ethanol for 2 min, rinsed thrice with sterilized distilled water, blot-dried for 3 min and transferred into potato dextrose agar (PDA) plates incubated at  $22\pm 2^{\circ}\text{C}$ , and newly produced sclerotia stored in 5ml screw cap glass tubes at  $4^{\circ}\text{C}$  (Atallah *et al.*, 2004).

### Preparation of culture filtrate

Two discs of mycelial agar plugs (5 mm diameter) from the margin of 7 days old culture of each isolate were placed into autoclaved 100 ml Richard's liquid media ( $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 2.5 g/l,  $\text{K}_2\text{HPO}_4$  5.0 g/l,  $\text{KNO}_3$  10.0 g/l,  $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ , 0.02g/l and sucrose 50 g/l in double distilled water sterilized) in 250 ml conical flasks and incubated at  $22\pm 1^{\circ}\text{C}$  on a rotary shaker at 100 rpm for 14 days. The mycelial mat was removed by filtering through a Whatman filter paper No.1 and heated at  $100^{\circ}\text{C}$  for 2 min to inactivate enzymes (Vidhysekaran *et al.*, 1970). Conical flasks with autoclaved 100 ml Richard's liquid media without mycelial agar plugs were kept as control.

### Variability in mycelial growth

Fresh mycelial weight was recorded. For dry mycelial weight, harvested mycelial mats were oven dried at  $60^{\circ}\text{C}$  for 72 hours.

### Effect of culture filtrate

Effect of *S. sclerotiorum* culture filtrates on germination and seedling vigour, one hundred seeds of *B. juncea* cv. Rohini were surface sterilized in 0.5% sodium hypochlorite ( $\text{NaClO}$ ) for 5 min., rinsed thrice in sterilized distilled water, placed on blotting paper in Petri plates, each watered with 10 ml of culture filtrate. Petri plates were sealed with parafilm, incubated in a growth chamber at  $25\pm 1^{\circ}\text{C}$  and 80% relative humidity, and percentage of seed germination and length of radical and plumule recorded 72 h and 7 days after incubation, respectively. The experimental design was completely randomized (CRD) with four replications.

### Results and Discussion

The growth rate of the isolates differed significantly. Maximum fresh mycelial weight was observed in SR 14 (13.9 g) followed by SR 08 (11.5 g), while

SR 17 (1.3 g) produced the maximum dry mycelial weight; SR 15 yielded the minimum fresh (0.96 g) and SR 22 the minimum dry mycelial weight (0.3 g), respectively (Table 1). Compared to the control treatments, culture filtrates of isolates SR 03 and SR 15 significantly reduced seed germination (53.5 and 53.3%, respectively) and caused maximum pre-emergence mortality (46.9 and 46.7%, respectively); seed germination (83.4%) and pre-emergence mortality (16.7%) were least affected by SR19 (Table 1). Post-emergence mortality was highest in SR 08 (80%) followed by SR 19 (76.7%); minimum post-emergence mortality was recorded in SR 02 (Table 1). Significant effect of culture filtrates was also observed on seedling vigour. For effect on lengths of radical and plumule, the culture filtrates of SR 08 and SR 14, and SR 05, 08, 09, 14, 15, 17, 20 and 22, respectively, proved to be most toxic (Table 1). Isolates SR 01 and SR 02 were least toxic in reducing lengths of radical and plumule, respectively (Table 1).

Variability in mycelial growth was observed among *S. sclerotiorum* isolates collected from different geographical regions in India. Morrall *et al.* (1972) also observed large variations in mycelial growth, numbers, shape, size and texture of sclerotia and other characteristics among 114 *Sclerotinia* spp. isolates collected from 23 hosts in Saskatchewan in Western Canada. Results of our study especially regarding variations in fresh and dry mycelial weight are also in agreement with the findings of Kohli *et al.* (1995) who reported remarkable phenotypic variability in mycelial growth rate, pigmentation, and amount of aerial mycelium in *S. sclerotiorum* populations, compared to the generally uniform appearance of isolates from canola in Canada and Norway. Willets and Wong (1980) also reported considerable variations in morphological characters among *S. Sclerotiorum*, *S. trifoliorum* and *S. minor* isolates.

The results of this study have shown that culture filtrates of different geographical isolates of *S. sclerotiorum* inhibit seed germination and seedling vigour including length of radical and plumule of Indian mustard. This shows that *S. sclerotiorum* produces toxic metabolites in the media in which they are grown. Pathogenic fungi may often damage their host plants by producing phytotoxins, which cause various symptoms including necrosis,

Table 1: Culture filtrate of different geographical isolates of *Sclerotinia Sclerotium* and their effect on seed germination and seedling vigour

Isolate no.	Fresh mycelial	Dry mycelial	Per cent seed germination	Pre-emergence mortality (%)	Post-emergence mortality (%)	Seedling vigour (mm)	
	wt (g)	wt (g)				Radicle	Plumule
SR01	4.21	0.58	70.0 (56.79)	30.0 (33.21)	46.7 (43.11)	7.33	9.67
SR02	1.65	0.53	63.3 (52.71)	36.7 (37.29)	26.7 (31.11)	5.33	10.67
SR03	3.18	0.78	53.5 (47.01)	46.9 (43.22)	33.3 (35.24)	3.33	7.00
SR04	5.40	0.89	76.7 (61.14)	23.3 (28.86)	46.7 (43.11)	2.12	3.67
SR05	8.78	0.67	73.3 (58.89)	26.7 (31.11)	63.3 (52.71)	1.00	0.00
SR06	6.42	1.16	63.3 (52.71)	36.7 (37.29)	50.0 (45.00)	2.67	3.67
SR07	3.57	0.95	63.3 (52.71)	36.7 (37.29)	43.3 (41.15)	1.50	3.33
SR08	11.51	0.74	80.0 (63.44)	20.0 (26.56)	80.0 (63.44)	0.10	0.10
SR09	4.88	0.82	70.0 (56.79)	30.0 (33.21)	66.7 (54.76)	0.33	0.00
SR10	4.25	1.26	70.0 (56.79)	30.0 (33.21)	56.7 (48.85)	5.67	4.00
SR11	8.97	0.57	73.3 (58.89)	26.7 (31.11)	60.0 (50.77)	3.67	4.00
SR12	4.68	0.75	66.7 (54.76)	33.3 (35.24)	43.3 (41.15)	2.00	1.33
SR13	5.23	0.85	76.7 (61.14)	23.3 (28.86)	63.3 (52.71)	4.67	2.67
SR14	13.87	0.55	73.3 (58.89)	26.7 (31.11)	73.3 (58.89)	0.10	0.10
SR15	0.96	0.52	53.3 (46.89)	46.7 (43.11)	33.3 (35.24)	1.33	0.00
SR16	5.61	0.75	73.3 (58.89)	26.7 (31.11)	63.3 (52.71)	1.00	0.33
SR17	9.00	1.27	66.7 (54.76)	33.3 (35.24)	63.3 (52.71)	0.67	0.00
SR18	2.64	0.98	56.7 (48.85)	43.3 (41.15)	46.7 (43.11)	4.00	2.67
SR19	3.30	0.78	83.3 (65.88)	16.7 (24.12)	76.7 (61.14)	1.00	0.33
SR20	7.92	0.39	66.7 (54.76)	33.3 (35.24)	63.3 (52.71)	0.67	0.00
SR21	2.91	0.73	70.0 (56.79)	30.0 (33.21)	43.3 (41.15)	8.00	2.00
SR22	7.14	0.33	70.0 (56.79)	30.0 (33.21)	66.7 (54.76)	0.33	0.00
SR23	3.76	0.65	73.3 (58.89)	26.7 (31.11)	60.0 (50.77)	3.67	6.33
SR24	4.33	1.04	76.7 (61.14)	23.3 (28.86)	73.3 (58.89)	0.33	0.33
SR25	3.45	0.89	63.3 (52.71)	36.7 (37.29)	33.3 (35.24)	5.33	4.33
Control (Un inoculated media)	0.00	0.00	90.0 (71.56)	10.0	5.31	60.42	14.21
Control (with distilled water)	0.00	0.00	96.7 (79.53)	3.33	0.00	111.0	52.33
CD (at 5%)	0.886	0.168	1.633	1.633	1.735	2.554	2.510
CV	10.179	13.699	14.168	11.261	9.946	12.835	13.468

\*Figures in parenthesis are angular transformed values

chlorosis, wilting, water soaking and eventually death of plants (Scheffer, 1983). Fungal filtrates from *Fusarium solani*, *F. oxysporum*, *Aspergillus niger*, *A. flavus*, *A. terreus* and *Alternaria alternata* were reported to reduce germination of soybean seeds (Ibraheem *et al.*, 1987). The culture filtrate of *Penicillium citrinum*, *Fusarium moniliforme* and *F. equiseti* also adversely affected the seed germination of fennel (Sharma and Sharma, 1983). The inhibition of radicle and plumule growth especially in wheat grown from seeds treated with *Fusarium head blight* fungus has been also reported (Gilbert and Tekauz, 1995). Metabolites from six different fungi also adversely affected seed germination and seedling growth in coriander (Pant, 2011).

## References

- Aggarwal RAK, KumRAK, Kumar A and Thakur HL. 1997. Effect of *Sclerotinia* rot on oil quality in low erucic acid cultivars of rapeseed. *Cruciferae Newslett* **19**: 103-104.
- Atallah ZK, Larget B, Chen X and Johnson DA. 2004. High genetic diversity, phenotypic uniformity, and evidence of outcrossing in *S. sclerotiorum* in the Columbia basin of Washington state. *Phytopathol* **94**: 737-742.
- Boland GJ and Hall R. 1994. Index of plant hosts of *Sclerotinia*. *Can J Plant Pathol* **16**: 93-108.
- Frame B, Kang-Fu Y, Christie BR and Paus KP. 1991. *In vitro* selection for resistance *Verticillium* wilt in alfalfa (*Medicago sativa* L.) using a fungal culture filtrate. *Physiol Mol Plant Pathol* **39**: 325-348.
- Gilbert J and Tekauz A. 1995. Effects of *Fusarium* head blight and seed treatment on germination, emergence and seedling vigour of spring wheat. *Canadian J Pl Pathol* **17**: 252-259.
- Griffin DH. 1981. Fungal Physiology. John Wiley and Sons, New York., pp: 383.
- Harlton CE, Lévesque CA and Punja ZK. 1995. Genetic diversity in *Sclerotium (Athelia) rolfsii* and related species. *Phytopathol* **85**: 1269-1281.
- Hilty TC and Lee HL. 1988. Hot acidified zinc sulphate as seed soaking agent for the control of crucifer black rot. *Plant Protec Bull* **30**: 245-248.
- Ibraheem SA, Okesha AM and Mhathem KT. 1987. Interrelationship between protein and oil content of soybean seed with some associated fungi. *J Agric Water Resources Res Plant Prod* **6**: 53-66.
- Kohli Y, Brunner LJ, Yoell H, Milgroom MG, Anderson JB, Morrall RAA and Kohn LM. 1995. Clonal dispersal and spatial mixing in populations of the plant-pathogenic fungus, *Sclerotinia sclerotiorum*. *Mol Ecol* **4**: 69-77.
- Madhosing C. 1995. Relative wilt-inducing capacity of the culture filtrates of isolates of *Fusarium oxysporum* f.sp. *radicis-lycopersici*, the tomato crown and root-rot pathogen. *J Phytopathol* **4**: 193-198.
- Morrall RAA, Duczek LJ and Sheard JW. 1972. Variation and correlations within and between morphology, pathogenicity, and pectolytic enzyme activity in *Sclerotinia* from Saskatchewan. *Can J Bot* **50**: 767-786.
- Okabe I, Morikawa C, Matsumoto N and Yokoyama K. 1998. Variation in *Sclerotium rolfsii* isolates in Japan. *Mycosci* **39**: 399-407.
- Pant R. 2011. Seed mycoflora of coriander and effect of some fungal metabolite on seed germination and seedling growth. *Asian J Exp Bio Sci* **2**: 127-130.
- Purdy LH. 1979. *Sclerotinia sclerotiorum*: history, diseases and symptomatology, host range, geographic distribution, and impact. *Phytopathol* **69**: 875-880.
- Scheffer RP. 1983. Toxins as chemical determinant of plant diseases. p. 1-40. In: *Toxins in plant pathogenesis*. (Eds.): JM Daly and BJ Darval, Academic Press, Sydney, 181.
- Sharma AK and Sharma KD. 1983. Effects of fungal metabolites on the germination of sweet fennel (*Foeniculum vulgare* Mill.) seeds. *Toxicology Letters* **17**: 81-84.
- Vidhyasekaran P, Subramanian CL and Govindaswamy CV. 1970. Production of toxins by seed-borne fungi and its role in paddy seed spoilage. *Indian Phytopath* **23**: 518-525.
- Willetts HJ and Wong AL. 1980. The biology of *S. sclerotiorum*, *S. trifoliorum* and *S. minor* with emphasis on specific nomenclature. *Bot Rev* **46**: 101-165.