



Variability in *Alternaria brassicae* and characterization of host response in Ethiopian mustard (*Brassica carinata* A. Braun)

Vedna Kumari¹, A Kumar³, HK Chaudhary¹, R Prasad², S Jambhulkar⁴ and S Sharma¹

¹Department of Crop Improvement

²Department of Agronomy

³Shivalik Agricultural Research & Extension Centre, Kangra
CSK Himachal Pradesh Krishi Vishvavidyalaya, Palampur-176062

⁴Bhabha Atomic Research Centre, Trombay, Mumbai, 400085

Corresponding author: drvedna@gmail.com

(Received: 13 Jan 2013; Revised: 16 April 2013; Accepted: 27 April 2013)

Abstract

The present investigation was undertaken to determine variability in virulence among four isolates of *Alternaria brassicae* (Berk.) Sacc., one each from *Brassica juncea*, *B. rapa*, *B. napus* and *B. carinata* on the basis of conidial morphology, cultural characteristics, sporulation intensity and length of incubation period. Results revealed that all parameters significantly affected virulence. The average length and breadth of the most virulent *B. napus* ($A.B_n$) isolate were 171.0 μm and 26.1 μm , compared to only 134.0 and 19.0 μm , respectively, of the least virulent *B. juncea* ($A.B_j$) isolate. Similarly, the horizontal and vertical conidial septations, diameters of both halo formation and concentric rings and sporulation intensity were significantly higher in the most virulent ($A.B_n$) than in the least virulent ($A.B_j$) isolates; incubation period was shortest (2 days) in the ($A.B_n$) isolate. Results of screening of 45 mutants of *B. carinata* against the most virulent $A.B_n$ isolate revealed longest incubation period of 4 to 5 days in the 17 mutants; 10 of the 17 mutants showed very poor sporulation intensity and smaller diameters of both halo formation and concentric rings. These 10 moderately resistant mutants viz; P_7 , (P_{10_2}), P_{11} , P_{74} in 50 kR, (P_{4_2}), (P_{5_2}), P_2 in 70 kR, (P_{4_2}) in 80 kR, P_{22} in 90 kR, and P_{13} in 100 kR were further field evaluated to confirm their reaction.

Key words: Ethiopian mustard, *Alternaria brassicae*, detached-leaf culture technique, virulence

Rapeseed-mustard are the most important *rabi* oilseed crops grown in India. In general, the crops have low average productivity due to the prevalence of various biotic and abiotic stresses. Diseases are among the major biotic stresses. *Alternaria* blight caused by *Alternaria brassicae* (Berk.) Sacc., is an important and a widespread destructive disease of rapeseed-mustard (Weiss, 1983). Depending on the severity of the disease, yield losses as high as 70 % have been reported in India (Kolte, 1985; Chahal, 1986; Saharan, 1991). Anamorph form of this pathogen shows great variability in morphology, physiology and pathogenicity. Several researchers have reported existence of variability based on morphology, sporulation, and growth and cultural characteristics (Gupta *et al.*, 1972; Saharan & Kadian, 1983; Awasthi and Kolte, 1989; Vishwanath and Kolte, 1997). Although, studies on pathogenic

variability are important for the development of pre-breeding populations (Meena *et al.*, 2010), information are lacking regarding differences in aggressiveness/virulence in the pathogenic population isolated from different *Brassica* species from Kangra region in Himachal Pradesh. The present investigation, therefore, was undertaken to determine the differences in virulence among the isolates from *Brassica juncea*, *B. rapa*, *B. napus* and *B. carinata* based on the morphology of conidia. The most virulent isolate discovered from these studies was subsequently used to screen mutants of *B. carinata* for their reaction against *A. brassicae*.

Materials and Methods

Alternaria brassicae-infected leaves, showing characteristic symptoms, were collected from plants of *B. juncea*, *B. rapa*, *B. napus* and *B. carinata*

grown in different fields of CSKHPKV, SAREC, Kangra, during *rabi*, 2010-11 season (Table 1). The pathogen was isolated, purified by single spore technique (Toussoun and Nelson, 1976), and maintained on PDA/ V₈ slants for further studies. Various isolates collected from different species were designated on the basis of host species *viz.*, *B. juncea* (*A.B_j*), *B. rapa* (*A.B_r*), *B. napus* (*A.B_n*) and *B. carinata* (*A.B_{car}*). The morphological characteristics of conidia of each isolate including length and breadth, number of septation (horizontal and transverse), and beak length were recorded in 15 days old cultures.

To determine the pathogenic variability in the laboratory, a detached leaf-culture technique was used (Bansal *et al.*, 1990). The samples of fresh and uniform-sized leaves of *B. carinata* variety 'Jayanti' were collected from the field, washed three times with distilled water and the lower epidermis from both sides of midrib was rubbed gently with the help of wet cotton. About 25 µl spore suspension (2000-2500 conidia/ml) of each isolate was injected on the punctured side of the leaves, and each leaf petiole was swabbed with wet cotton. The leaves were kept inside the wet chambers at 25±1°C for 48-72 hrs. The observations on incubation period, halo formation, concentric ring size, and sporulation were recorded periodically to assess the virulence of four isolates. The most virulent isolate was then used to screen 45 putative mutants of *B. carinata* for their reaction against *A. brassicae*.

Table 1: Name of isolates and their codes

Host species	Isolate code
Brassica juncea	A.B _j
B. rapa	A.B _r
B. napus	A.B _n
<i>B. carinata</i>	<i>A.B_{car}</i>

Results and Discussion

Conidial characteristics of the isolates:

Morphological observations revealed that the isolates differed in their conidial size (Table 2). Amongst the isolates, the average conidial length varied from 134.0 to 171.0 µm. The maximum

(358 µm) and minimum (134 µm) length, respectively, were observed in isolate *A.B_n* and *A.B_j*, although, both were statistically at par with each other, the conidial length varied from 80 to 358 µm. The average breadth of conidia varied from 19.0 to 26.1 µm. The isolate *A.B_n* exhibited thickest conidium (26.1 µm) while the isolate *A.B_{car}* was the thinnest (19.0 µm), followed by *A.B_j* (19.1 µm). The isolate *A.B_n* showed maximum range of conidial breadth that varied from 8 to 38 µm. Between *brassica* species, isolates did not differ greatly in the average number of septations. Between isolates, the horizontal septations ranged from 4 to 18, and vertical from 0 to 9. Our observations support the report by Kumar *et al.* (2003) where they also observed very little variation in both horizontal (3 to 12) and vertical septation (0 to 6). A significant variation in the beak length between isolates was observed. Although, the average beak length varied from 25.0 to 46.0 µm, it was significantly longer in isolate *A.B_n* and smaller in *A.B_j*. The longest and smallest beak lengths were observed in isolate *A.B_{car}* (188 µm) and isolate *A.B_j* (68 µm), respectively. Similar variations in length and breadth of conidia have also been reported previously (Ellis, 1971; Holliday, 1980; Kumar *et al.* 2003; Singh *et al.* 2009; Meena *et al.* 2012). Kumar *et al.* (2003) also reported variations in the average length (118.6 to 194.5 µm) and breadth (14.5 to 22.7 µm) in *A. brassicae* conidia. Singh *et al.* (2009) also observed variability in conidial length (117.0 to 192.0 µm), breadth (14.0 to 24.0 µm), conidial beak length (42.0 to 116.0 µm), and number of horizontal and longitudinal septa. High level of variability in conidial length, width, beak length, and number of septa *in-vitro* produced conidia of different isolates was also observed by Meena *et al.* (2012). Slight variations in the conidial size between studies may be attributed to the physical conditions of the medium (Plate 1). Variations in the conidial size in the present and several earlier investigations suggest existence of variability in this pathogen.

Plate 1: *Alternaria brassicae* conidia from four *Brassica* species.




(1) *A.B_j* spores /conidia



(2) *A.B_r* spores/conidia



(3) *A.B_n* spores /conidia



(4) *A.B_{car}* spores/conidia

- (1) *A.B_j* – Spores/conidia of *Alternaria brassicae* isolated from *Brassica juncea*
 (2) *A.B_r* – Spores/conidia of *Alternaria brassicae* isolated from *Brassica rapa*
 (3) *A.B_n* – Spores/conidia of *Alternaria brassicae* isolated from *Brassica napus*
 (4) *A.B_{car}* – Spores/conidia of *Alternaria brassicae* isolated from *Brassica carinata*

Pathogenicity of *Alternaria brassicae* isolates on *Brassica carinata*:

The pathogenic variability of four *Alternaria brassicae* isolates was observed on *Brassica carinata* on the basis of parameters including incubation period, halo formation, concentric ring size and sporulation intensity (Table 3, Plate 2).

It was observed that the length of incubation period, size of concentric ring, halo formation and sporulation intensity were directly related to virulence of the isolates. The virulent isolate, thus,

causes higher disease intensity by inducing bigger halo formation, larger concentric rings, higher sporulation, and significantly shorter mean incubation period. The most virulent isolate *A.B_n* recorded shortest incubation period, largest diameter of concentric ring and halo formation and highest sporulation intensity on *Brassica carinata*. Virulence was also related to the morphology of conidia. The most virulent *A.B_n* isolate produced significantly larger sized conidia compared to the least virulent *A.B_j* isolate, thereby indicating a direct relation between virulence and size of conidia. The

Table 2: Variation in conidial size of *Alternaria brassicae* isolates obtained from different *Brassica* species

Isolates	<u>Length (µm)</u>		<u>Breadth (µm)</u>		<u>Septation (No.)</u>				<u>Beak length (µm)</u>	
	A	R	A	R	<u>Horizontal</u>		<u>Vertical</u>		A	R
					A	R	A	R		
<i>A.B_j</i>	134.0	80-200	19.1	13-32	8	4-14	1	0-4	25.0	10-68
<i>A.B_r</i>	154.0	88-288	21.0	16-35	9	4-16	3	0-7	40.0	18-122
<i>A.B_n</i>	171.0	105-358	26.1	8-38	10	5-18	2	0-9	46.0	16-150
<i>A.B_{car}</i>	142.0	85-250	19.0	13-28	9	4-15	1	0-5	43.0	13-188
Mean	178.6		20.0						43.6	
CD (5%)	41.4		NS						10.3	
CV (%)	15.1		12.4						15.4	

A = Average, R= Range

Table 3: Pathogenicity of four isolates of *Alternaria brassicae* on *Brassica carinata* through detached leaf-culture technique

<i>A. brassicae</i> isolates	Mean incubation period	Halo Formation (days)	Concentric ring size (mm)	Sporulation intensity (mm)	Virulence categories
<i>A.B_n</i>	2.0	120	60	much dark	highly virulent
<i>A.B_r</i>	3.0	90	56	dark	moderate virulent
<i>A.B_{car}</i>	3.7	75	50	medium light	moderate virulent
<i>A.B_j</i>	3.3	50	40	very light	least virulent

Plate 2: Detached leaf- culture technique – Test for the pathogenicity of four isolates

**Reactions of different *Brassica* species isolates on *Brassica carinata*.**(A) *A.B_n* (B) *A.B_r* (C) *A.B_{car}* (D) *A.B_j*

findings of the present investigation clearly demonstrated existence of pathogenic variability in *Alternaria brassicae* isolated from different *Brassica* species. Wide variation in virulence among *Alternaria brassicae* isolates from rapeseed-mustard have also been observed by Saharan and

Kadian (1983). In the present investigation, the most virulent isolate was obtained from *B. napus* thus, partially supporting the observations of Bains and Tewari (1987) where they reported existence of resistance in *Brassica napus*, *Brassica carinata* and *Brassica rapa* against *Alternaria* blight. This

Table 4: Response of 45 putative mutants of *Brassica carinata* to the most virulent isolate of *A. brassicae* (*A.B_n*) through detached leaf-culture technique

<i>B. carinata</i> mutants	Dose	Mean incubation period (days)	Mean halo diameter (mm)	Mean concentric ring size (mm)	Sporulation intensity	Category
P ₇	50 kR	4.5	40	38	very less	MR
(P ₁₀) ₂	50 kR	4.0	37	37	very less	MR
P ₁₀	50 kR	3.0	68	60	dark	HS
P ₄₆	50 kR	3.9	64	56	dark	S
(P ₈) ₂	50 kR	4.2	42	41	medium	MS
P ₁₁	50 kR	3.7	40	40	very less	MR
P ₇₄	50 kR	4.0	45	42	very less	MR
(P ₁₁) ₂	50 kR	4.4	48	40	medium	MS
P ₅₆	50 kR	3.8	56	49	dark	S
(P ₃) ₂	50 kR	3.6	73	61	dark	S
P ₄₇	50 kR	4.2	57	50	dark	S
P ₂₆	60 kR	3.8	65	55	medium	MS
P ₃₉	60 kR	3.5	56	51	medium	MS
P ₃₈	60 kR	3.0	75	68	very dark	HS
(P ₁) ₂	60 kR	3.8	63	58	sparsely	MS
(P ₉) ₂	60 kR	3.9	65	55	dark	S
(P ₄) ₂	70 kR	4.4	45	36	very less	MR
(P ₅) ₂	70 kR	4.5	48	45	very less	MR
P ₆	70 kR	3.7	78	70	dark	S
P ₂₂	70 kR	4.0	58	50	medium	MS
(P ₂) ₂	70 kR	4.2	50	47	medium	MS
P ₂	70 kR	4.0	47	43	very less	MR
(P ₄) ₂	80 kR	4.4	45	36	very less	MR
P ₁₅	90 kR	3.0	72	70	dark	HS
P ₂	90 kR	4.0	66	50	medium	MS
P ₂₂	90 kR	4.5	35	35	very less	MR
(P ₂) ₂	90 kR	4.0	50	38	medium	MS
P ₂₂	100 kR	3.9	58	50	medium	MS
P ₄	100 kR	3.7	56	49	medium	MS
(P ₁) ₂	100 kR	3.8	60	55	medium	MS
P ₁₃	100 kR	4.3	42	40	sparsely	MR
(P ₂) ₂	110 kR	3.5	68	54	medium	S
(P ₂) ₂	0.3% EMS-PS	3.0	75	70	very dark	HS

P ₃₄	0.3% EMS-PS	3.0	72	61	very dark	HS
P ₃₁	0.3% EMS-PS	3.0	74	60	medium	S
P ₁₃	0.3% EMS-PS	3.7	60	52	medium	S
(P ₉) ₂	0.3% EMS-PS	3.1	68	59	dark	S
(P ₁₇) ₂	0.3% EMS-PS	4.0	50	45	medium	MS
P ₂₇	0.3% EMS-PS	3.0	56	50	medium	S
P ₁₄	0.3% EMS-PS	3.5	54	50	very less	MS
P ₁₈	0.3% EMS-PS	3.2	58	52	sparsely	MS
P ₅	0.3% EMS-PS	3.0	69	65	very dark	S
(P ₂₃) ₂	0.3% EMS-PS	3.0	74	70	very dark	HS
(P ₁) ₂	0.4% EMS-PS	3.1	71	67	dark	HS
P ₈	0.5% EMS-PS	3.5	62	55	medium	S

MR-Moderately Resistant; MS-Moderately Susceptible; S-Susceptible, HS-Highly susceptible; EMS-PS-Ethyl Methane Sulphonate with pre-soaking

virulent isolate we obtained can now be utilized in screening rapeseed-mustard germplasm and advanced breeding lines for resistance against *Alternaria brassicae*.

Response of *Brassica carinata* mutants to the most virulent *Alternaria brassicae* isolate:

The most virulent *B. napus* isolate *A.B_n*, was used to screen 45 putative mutants of *Brassica carinata*. The longest incubation period of 4-5 days was recorded in 17 mutants, 10 of the 17 mutants showed very poor sporulation intensity and smaller diameters of both halo and concentric rings (Table 4). Because of their moderate resistance to *A. brassicae*, these 10 mutants viz; P₇, (P₁₀)₂, P₁₁ and P₇₄ in 50 kR, (P₄)₂, (P₅)₂ and P₂ in 70 kR, (P₄)₂ in 80 kR, P₂₂ in 90 kR and P₁₃ in 100 kR, were further evaluated under natural field conditions at Kangra.

Acknowledgement

The senior author is grateful to the Bhabha Atomic Research Centre (BARC), Trombay, Mumbai, for providing financial assistance in the form of an ad-hoc research project for the present study.

References

Awasthi, RP and Kolte, SJ. 1989. Variability in *Alternaria Brassicae* affecting rapeseed and mustard. *Indian Phytopath* **42**: 275.

Bains, PS and Tewari, JP. 1987. Purification, chemical characterization and host specificity of the toxin produced by *Alternaria brassicae*. *Physio Mol Plant Pathol* **30**: 259-271.

Bansal, VK, Séguin- Swartz, G, Rakow G and Petrie, GA. 1990. Reaction of *Brassica* species to infection by *Alternaria brassicae*. *Can J Plant Sci* **70**: 1159-1162.

Chahal, AS. 1986. Losses and chemical control of *Alternaria* in rapeseed-mustard in Punjab. *Plant Disease Res* **1**: 46-50.

Ellis, MB. 1971. *Dematiaceous hyphomycetes*. Commonwealth Mycological Institute, England. No. 608: 464-497pp.

Gupta, RBL, Desai, BG and Pathak, VN. 1972. Effect of light on growth and sporulation of *Alternaria brassicae* (Berk.) Sacc. *Phytopathol Mediterr* **11**: 61-62.

Holliday, P. 1980. *Fungal Diseases of Tropical Crops*. Cambridge Univ. press, London. 6-19pp.

Kolte, SJ. 1985. *Diseases of Annual Edible Oilseed Crops*. Vol. II, Rapeseed-Mustard and Sesame Diseases. CRC Press, Florida. 135pp.

Kumar, S, Sangwan, MS, Mehta, N and Kumar, R. 2003. Pathogenic diversity in isolates of *Alternaria brassicae* infecting rapeseed and Mustard. *J Mycol Pl Pathol* **33**: 59-64.

Meena, PD, Awasthi, RP, Chattopadhyay, C, Kolte, SJ and Kumar, A. 2010. *Alternaria* blight: a chronic disease in rapeseed-mustard. *J Oilseed Brassica* **1**: 1-11.

Meena, PD, Rani, A, Meena, R, Sharma, P, Gupta, R and Chowdappa, P. 2012. Aggressiveness, diversity and distribution *Alternaria brassicae*

- isolates infecting oilseed Brassica in India. *African J Microbiol Res* 6 (24): 5249-5258.
- Saharan, GS. 1991. Assessment of losses, epidemiology and management of black spot disease of rapeseed-mustard. Proceedings GCIRC 8th Intern. Rapeseed Congr. (July 9-11), Saskatoon, Canada, Vol. II: 465-470 84pp (Abstr.)
- Saharan, GS and Kadian, AK. 1983. Physiological specialization in *Alternaria brassicae*. *Cruciferae Newsletter* 8: 32-33.
- Singh, D, Singh, R, Singh, H, Yadav, RC, Yadav, N, Barbetti, M and Salisbury, P. 2009. Cultural and morphological variability in *Alternaria brassicae* isolates of Indian mustard, *Brassica juncea* L. Czern & Coss. *J Oilseed Res.* 26: 134-137.
- Toussoun, TA and Nelson, PE. 1976. A pictorial guide to the identification of *Fusarium* spp. According to the Taxonomic system of Synder and Hanson. University Park, USA, 43pp.
- Vishwanath and Kolte, SJ. 1997. Variability in *Alternaria brassicae*: Response to host genotypes, toxin production and fungicides. *Indian Phytopath* 50: 373-381.
- Weiss, E.A. 1983. *Oilseed Crops*. Longman Inc. New York.

